

OPTIMAL HOMOTOPY ASYMPTOTIC METHOD FOR FIRST-ORDER CONFORMABLE FRACTIONAL DIFFERENTIAL EQUATIONS

M. ILIE, J. BIAZAR, Z. AYATI

ABSTRACT. In this article, optimal homotopy asymptotic method is presented to solve linear and nonlinear first-order conformable fractional differential equations that, is named conformable fractional optimal homotopy asymptotic method. So conformable fractional relaxation-oscillation and Riccati differential equations as a linear and nonlinear fractional differential equations, are solved by the proposed approach, respectively. The results obtained demonstrate the efficiency of the declared method for fractional equations.

1. INTRODUCTION

Many phenomena in our real world are described by fractional differential equations [3-14]. Fractional differential equations are often seeming perplexing to solve. Therefore, finding comprehensive methods for solving them sounds of high importance. Although having the exact solution of fractional equations in analyzing the phenomena is essential, there are many fractional differential equations, which cannot be solved exactly. Due to this fact, finding a desired approximate solutions of fractional differential equations is clearly vital. In recent years, many effective methods have been proposed for finding approximate solution to fractional differential equations, such as Adomian decomposition method [14, 15], homotopy perturbation method [16-19], homotopy analysis method [20], optimal homotopy asymptotic method [21-23], variational iteration method [24], generalized differential transform method [25], finite difference method [26], semi-discrete scheme and Chebyshev collocation method [27], wavelet operational [28, 29] and other methods [30-36]. In this paper, optimal homotopy asymptotic method is utilized to obtain an approximate solution of linear and nonlinear conformable fractional differential equations. The relaxation-oscillation and Riccati fractional differential equations, such as a linear and a nonlinear fractional equations are solved, respectively. The organization of this paper is as follows: In Section 2, the basic definitions like conformable fractional derivative and integral, is described. In Section 3, the conformable fractional

2010 *Mathematics Subject Classification.* 34A08, 34A12, 34A30.

Key words and phrases. Conformable fractional Riccati differential equations, Conformable fractional relaxation-oscillation differential equations, Optimal homotopy asymptotic method, Conformable fractional derivative.

Submitted November 17, 2017.

homotopy asymptotic method for linear and nonlinear fractional differential equations, are presented. In Section 4, conformable fractional relaxation-oscillation and Riccati equations, as illustrative examples, by means of the proposed approach are solved. Finally, conclusions are given in Section 5.

2. BASIC DEFINITIONS

The purpose of this section is to recall some preliminaries of the proposed method.

2.1. Conformable fractional derivative (CFD). Given a function $f : [0, \infty) \rightarrow \mathbb{R}$. The conformable fractional derivative of f , of order α , is defined by

$$T_\alpha(f)(x) = \lim_{\varepsilon \rightarrow 0} \frac{f(x + \varepsilon x^{1-\alpha}) - f(x)}{\varepsilon}, \quad (2.1)$$

for all $x > 0$, $\alpha \in (0, 1]$. If f is α -differentiable in some interval $(0, a)$, let's define $T_\alpha(f)(0) = \lim_{x \rightarrow 0^+} T_\alpha(f)(x)$, provided that $\lim_{x \rightarrow 0^+} T_\alpha(f)(x)$ exists. If the conformable derivative of f of order α exists, then we simply say that f is α -differentiable (see [1, 2]).

One can easily show that satisfies all the following properties (see [1]):

Let $a \in (0, 1]$ and f, g be a -differentiable at a point $x > 0$, then

- A. For $a, b \in \mathbb{R}$ $T_\alpha(af + bg) = aT_\alpha(f) + bT_\alpha(g)$,
- B. For all $p \in \mathbb{R}$ $T_\alpha(x^p) = px^{p-\alpha}$,
- C. For all constant functions $f(x) = \lambda$, $T_\alpha(\lambda) = 0$,
- D. $T_\alpha(f \cdot g) = g \cdot T_\alpha(f) + f \cdot T_\alpha(g)$,
- E. $T_\alpha\left(\frac{f}{g}\right) = \frac{g \cdot T_\alpha(f) - f \cdot T_\alpha(g)}{g^2}$,
- F. $T_\alpha(f) = x^{1-\alpha} \frac{df}{dx}$.

If $\alpha \in (n, n + 1]$ and is n -differentiable at $x > 0$, then the conformable fractional derivative of order α is defined as follows

$$T_\alpha(f)(x) = \lim_{\varepsilon \rightarrow 0} \frac{f^{([\alpha]-1)}(x + \varepsilon x^{([\alpha]-\alpha)}) - f^{([\alpha]-1)}(x)}{\varepsilon}, \quad (2.2)$$

where $[\alpha]$ is the smallest integer greater than or equal to α . When is $(n + 1)$ -differentiable at $x > 0$, as a consequence of (2.2), one can have (see [1])

$$T_\alpha(f)(x) = x^{[\alpha]-\alpha} \frac{d^{([\alpha])}f}{dx^{([\alpha])}}.$$

2.2. Conformable fractional integral. Given a function $f : [a, \infty) \rightarrow \mathbb{R}$, $a \geq 0$. The conformable fractional integral of f , is defined by

$$I_\alpha^a(f)(x) = \int_a^x \frac{f(t)}{t^{1-\alpha}} dt, \quad (2.3)$$

where the integral is the usual Riemann improper integral, and $\alpha \in (0, 1)$ (see [1, 2]). For the sake of simplicity, let's consider $I_\alpha^0(f)(x) = I_\alpha(f)(x)$. One of the most useful results is the following statement (see [1]):

For all $x \geq a$ and any continuous function in the domain of I_α^a , we have $T_\alpha(I_\alpha^a f(x)) = f(x)$.

3. Conformable fractional optimal homotopy asymptotic method

Consider conformable fractional functional equation with initial condition, as the following

$$T_\alpha(u) = \mathcal{N}(u) + f(t), u(0) = \beta, \tag{3.1}$$

where \mathcal{N} is a functional operator, $f(t)$ is a known function, and β is a constant. The Eq. (3.1) is changed into conformable fractional Riccati differential equation, when $\mathcal{N}(u) = g(t)u(t) + h(t)u^2(t)$, where g, h are given functions.

If $\mathcal{N}(u) = Bu(t)$, then Eq. (3.1) is converted into conformable fractional relaxation-oscillation differential equation, where B is a constant.

According to optimal homotopy asymptotic technique, a homotopy $v(t, p) : \Omega \times [0, 1] \rightarrow \mathbb{R}$ can be constructed as the following

$$(1 - p) [T_\alpha(v(t, p)) - f(t)] - H(p) [T_\alpha(v(t, p)) - \mathcal{N}(v(t, p)) - f(t)] = 0. \tag{3.2}$$

Where $p \in [0, 1]$ is an embedding parameter, $H(p)$ is a nonzero auxiliary function for $p \neq 0$ and $H(0) = 0$, $v(t, p)$ is an unknown function. By substituting $p = 0$ and 1 in equation (3.2), we have and respectively. Thus as p is changing from zero to unity, the solution $v(t, p)$ varies continuously from $u_0(t)$ to an exact solution $u(t)$. By substituting $p = 0$ in Eq. (3.2) the initial approximation $u_0(t) = v(t, 0)$ is obtained as the solution of equation, the following

$$T_\alpha(u_0(t)) - f(t) = 0, u(0) = \beta. \tag{3.3}$$

The auxiliary function $H(p)$ can be chose as the following

$$H(p) = c_1p + c_2p^2 + c_3p^3 + \dots, \tag{3.4}$$

where c_1, c_2, c_3, \dots are parameters, that will be determined later. Expanding $v(t, p, c_1, c_2, \dots)$ in a Taylor series of p is

$$v(t, p, c_1, c_2, \dots) = u_0(t) + \sum_{i=1}^{\infty} u_i(t, c_1, c_2, \dots, c_i) p^i. \tag{3.5}$$

Substituting Eqs. (3.4) and (3.5) into equation (3.2) and setting to zero the coefficient of like powers of p , then the zero order deformation equation is obtained as given in Eq. (3.3), and the other order deformation equations are given as the following

$$\begin{aligned} p^1; & T_\alpha(u_1(t, c_1)) - T_\alpha(u_0(t)) + f(t) - c_1 [T_\alpha(u_0(t)) - \mathcal{N}(u_0(t)) - f(t)] = 0, u_1(0) = 0, \\ p^2; & T_\alpha(u_2(t, c_1, c_2)) - T_\alpha(u_1(t, c_1)) - c_2 [T_\alpha(u_0(t)) - \mathcal{N}(u_0(t)) - f(t)] \\ & - c_1 [T_\alpha(u_1(t, c_1)) - u_1(t, c_1) \frac{\partial \mathcal{N}}{\partial u_0}(u_0(t))] = 0, u_2(0) = 0, \\ p^3; & T_\alpha(u_3(t, c_1, c_2, c_3)) - T_\alpha(u_2(t, c_1, c_2)) - c_3 [T_\alpha(u_0(t)) - \mathcal{N}(u_0(t)) - f(t)] \\ & - c_2 [T_\alpha(u_1(t, c_1)) - u_1(t, c_1) \frac{\partial \mathcal{N}}{\partial u_0}(u_0(t))] \\ -\frac{1}{2}c_1 & [2T_\alpha(u_2(t, c_1, c_2)) - u_1^2(t, c_1) \frac{\partial^2 \mathcal{N}}{\partial u_0^2}(u_0(t)) - 2u_2(t, c_1, c_2) \frac{\partial \mathcal{N}}{\partial u_0}(u_0(t))] = 0, u_3(0) = 0, \\ & \vdots \end{aligned} \tag{3.6}$$

It should be noted that u_1, u_2, u_3, \dots are directed by linear equations (3.6), which can be easily solved. The convergence of the series given in Eq. (3.5) depends upon

the auxiliary parameters c_i for $i \geq 1$ If it converges at $p = 1$ we have,

$$u(t, c_1, c_2, \dots) = u_0(t) + \sum_{i=1}^{\infty} u_i(t, c_1, c_2, \dots, c_i). \quad (3.7)$$

Generally, the m th order approximate solution of Eq. (3.1), can be denoted as the following

$$u^m(t, c_1, c_2, \dots, c_m) = u_0(t) + \sum_{i=1}^m u_i(t, c_1, c_2, \dots, c_i). \quad (3.8)$$

By substitution of Eq. (3.8) into Eq. (3.1), the residual error can be expressed as follows

$$R(t, c_1, c_2, \dots, c_m) = T_\alpha(u^m(t, c_1, c_2, \dots, c_m)) - \mathcal{N}(u^m(t, c_1, c_2, \dots, c_m)) - f(t). \quad (3.9)$$

When $R(t, c_1, c_2, \dots, c_m) = 0$, results that $u^m(t, c_1, c_2, \dots, c_m)$ is an exact solution. Such a case does not usually occur for nonlinear problems. In these cases we can apply least square approach:

$$J_m(c_1, c_2, \dots, c_m) = \int_a^b R^2(t, c_1, c_2, \dots, c_m) dt, \quad (3.10)$$

where the values a, b depend on the given problem. The unknown convergence control parameters $c_1, c_2, c_3, \dots, c_m$ can be optimally identified from the following conditions

$$\frac{\partial J_m}{\partial c_i} = 0, i = 1, 2, \dots, m. \quad (3.11)$$

It is interesting to point out that when these parameters are determined, the m th order approximate solution given by Eq. (3.8) can be constructed.

4. EXAMPLES

In this section, to illustrate the proposed approach, conformable fractional relaxation-oscillation and Riccati differential equations will be solved.

Example 4.1. Consider the following fractional relaxation-oscillation differential equation with initial value

$$T_{\frac{3}{2}} u(t) = -u(t), 0 \leq t \leq 1, u(0) = 1, \quad (4.1)$$

where $u(x) = \exp\left(-\frac{2}{3}t^{\frac{3}{2}}\right)$.

According to the proposed conformable fractional optimal homotopy asymptotic method, we have

$$(1-p) \left[T_{\frac{3}{2}}(\nu(t, p)) \right] - H(p) \left[T_{\frac{3}{2}}(\nu(t, p)) + \nu(t, p) \right] = 0. \quad (4.2)$$

Substituting Eqs. (3.4) and (3.5) into Eq. (4.2) and setting to zero the coefficient of the same powers of p , we derive

$$p^0; T_{\frac{3}{2}}(u_0(t)) = 0, u_0(0) = 1,$$

$$p^1; T_{\frac{3}{2}}(u_1(t, c_1)) - T_{\frac{3}{2}}(u_0(t)) - c_1 \left[T_{\frac{3}{2}}(u_0(t)) + u_0(t) \right] = 0, u_1(0) = 0, \quad (4.3)$$

$$\begin{aligned}
 p^2; \mathbb{T}_{\frac{3}{2}} (u_2 (t, c_1, c_2)) - \mathbb{T}_{\frac{3}{2}} (u_1 (t, c_1)) - c_2 \left[\mathbb{T}_{\frac{3}{2}} (u_0 (t)) + u_0 (t) \right] \\
 - c_1 \left[\mathbb{T}_{\frac{3}{2}} (u_1 (t, c_1)) + u_1 (t, c_1) \right] = 0, \quad u_2 (0) = 0, \\
 \vdots
 \end{aligned}$$

Corresponding solution of this system of conformable differential equations (4.3), are

$$\begin{aligned}
 u_0 (t) &= 1, \\
 u_1 (t, c_1) &= \frac{2}{3} c_1 t^{\frac{3}{2}}, \\
 u_2 (t, c_1, c_2) &= \frac{2}{3} c_1 t^{\frac{3}{2}} + \frac{2}{3} c_2 t^{\frac{3}{2}} + \frac{2}{3} c_1^2 t^{\frac{3}{2}} + \frac{2}{9} c_1^2 t^3, \\
 &\vdots
 \end{aligned}$$

Therefore, two-terms approximation to the solution of Eq. (4.1), can be obtained as follows,

$$\begin{aligned}
 u^2 (t, c_1, c_2) &= 1 + 1.3333333333 c_1 t^{\frac{3}{2}} + 0.6666666667 c_2 t^{\frac{3}{2}} \\
 &\quad + 0.6666666667 c_1^2 t^{\frac{3}{2}} + 0.2222222222 c_1^2 t^3.
 \end{aligned} \tag{4.4}$$

For calculating unknown auxiliary constants c_1 and c_2 in $u^2(t, c_1, c_2)$ given in Eq. (4.4), we apply the procedure mentioned in (3.9) up to (3.11), we one obtains

$$c_1 = 0.8647018968, \quad c_2 = -3.456662450.$$

By considering these values, in (4.4), two-order approximate solution of Eq. (4.1) reads to

$$u^2(t) = 1 - 0.653032857t^{\frac{3}{2}} + 0.1661576378t^3.$$

In Figure [4.1], the exact and approximate solutions of fractional relaxation-oscillation (4.1) equation are plotted.

Example 4.2. Consider the following linear fractional relaxation-oscillation differential equation with initial value

$$\mathbb{T}_\alpha u(t) = 1 + u(t), \quad 0 \leq t, \quad 0 < \alpha \leq 1, \quad u(0) = 0. \tag{4.5}$$

The exact solution of Eq. (4.4), is $u(t) = \exp\left(\frac{1}{\alpha}t^\alpha\right) - 1$.

According to the proposed conformable fractional optimal homotopy asymptotic method, we have

$$(1 - p) [\mathbb{T}_\alpha (v(t, p)) - 1] - H(p) [\mathbb{T}_\alpha (\nu(t, p)) - \nu(t, p) - 1] = 0. \tag{4.6}$$

Substituting Eqs. (3.4) and (3.5) into Eq. (4.6) and setting to zero the coefficient of like powers of p , we obtain

$$\begin{aligned}
 p^0; \mathbb{T}_\alpha (u_0 (t)) - 1 &= 0, \quad u_0 (0) = 0, \\
 p^1; \mathbb{T}_\alpha (u_1 (t, c_1)) - \mathbb{T}_\alpha (u_0 (t)) + 1 - c_1 [\mathbb{T}_\alpha (u_0 (t)) - u_0 (t) - 1] &= 0, \quad u_1 (0) = 0, \\
 p^2; \mathbb{T}_\alpha (u_2 (t, c_1, c_2)) - \mathbb{T}_\alpha (u_1 (t, c_1)) - c_2 [\mathbb{T}_\alpha (u_0 (t)) - u_0 (t) - 1] \\
 - c_1 [\mathbb{T}_\alpha (u_1 (t, c_1)) - u_1 (t, c_1)] &= 0, \quad u_2 (0) = 0, \\
 p^3; \mathbb{T}_\alpha (u_3 (t, c_1, c_2, c_3)) - \mathbb{T}_\alpha (u_2 (t, c_1, c_2)) - c_3 [\mathbb{T}_\alpha (u_0 (t)) - u_0 (t) - 1] \\
 - c_2 [\mathbb{T}_\alpha (u_1 (t, c_1)) - u_1 (t, c_1)] - c_1 [\mathbb{T}_\alpha (u_2 (t, c_1, c_2)) - u_2 (t, c_1, c_2)] &= 0, \quad u_3 (0) = 0, \\
 &\vdots
 \end{aligned} \tag{4.7}$$

Matching solution of conformable equations (4.7), are

$$\begin{aligned} u_0(t) &= \frac{1}{\alpha} t^\alpha, \\ u_1(t, c_1) &= -\frac{1}{2} c_1 \left(\frac{1}{\alpha} t^\alpha\right)^2, \\ u_2(t, c_1, c_2) &= -\frac{1}{2} c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^2 - \frac{1}{2} c_1 \left(\frac{1}{\alpha} t^\alpha\right)^2 + \frac{1}{6} c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^3 - \frac{1}{2} c_2 \left(\frac{1}{\alpha} t^\alpha\right)^2, \\ u_3(t, c_1, c_2, c_3) &= -\frac{1}{2} c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^2 + \frac{1}{3} c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^3 - c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^2 - c_1 c_2 \left(\frac{1}{\alpha} t^\alpha\right)^2 \\ &\quad - \frac{1}{24} c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^4 + \frac{1}{3} c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^3 + \frac{1}{3} c_1 c_2 \left(\frac{1}{\alpha} t^\alpha\right)^3 - \frac{1}{2} c_2 \left(\frac{1}{\alpha} t^\alpha\right)^2 - \frac{1}{2} c_3 \left(\frac{1}{\alpha} t^\alpha\right)^2, \\ &\quad \vdots \end{aligned}$$

Thus, third-terms approximation to the solution of Eq. (4.5), will be obtained as the following

$$\begin{aligned} u^3(t, c_1, c_2, c_3) &= \left(\frac{1}{\alpha} t^\alpha\right) - 1.5c_1 \left(\frac{1}{\alpha} t^\alpha\right)^2 - 1.5c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^2 + 0.5c_1^2 \left(\frac{1}{\alpha} t^\alpha\right)^3 - c_2 \left(\frac{1}{\alpha} t^\alpha\right)^2 \\ &\quad - 0.5000000001c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^2 + 0.3333333334c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^3 - c_1 c_2 \left(\frac{1}{\alpha} t^\alpha\right)^2 \\ &\quad - 0.04166666668c_1^3 \left(\frac{1}{\alpha} t^\alpha\right)^4 + 0.3333333334c_1 c_2 \left(\frac{1}{\alpha} t^\alpha\right)^3 - 0.5000000001c_3 \left(\frac{1}{\alpha} t^\alpha\right)^2. \end{aligned} \quad (4.8)$$

Table 4.1, shows the optimal values of the convergence control constants c_1 , c_2 and c_3 in $u^3(t, c_1, c_2, c_3)$ given in Eq. (4.8) for different values of which can be obtained using the procedure mentioned in (3.9) up to (3.11). In Figures 4.2, the an exact and approximate solutions of fractional Relaxation-Oscillation equation for $\alpha = 0.4, 0.6, 0.8$, and 1.0 are plotted.

Example 4.3. Consider the following conformable fractional Riccati differential equation with initial value

$$\mathbb{T}_\alpha u(t) = 1 - u^2(t), \quad 0 \leq t \leq 1, \quad 0 < \alpha \leq 1, \quad u(0) = 0, \quad (4.9)$$

where $u(t) = \frac{\exp(\frac{2}{\alpha} t^\alpha) - 1}{\exp(\frac{2}{\alpha} t^\alpha) + 1}$.

Consistent with the conformable fractional homotopy asymptotic method, we obtain

$$(1-p) [\mathbb{T}_\alpha(v(t,p)) - 1] - H(p) [\mathbb{T}_\alpha(v(t,p)) + v^2(t,p) - 1] = 0. \quad (4.10)$$

Substituting Eqs. (3.4) and (3.5) into Eq. (4.10) and setting to zero the coefficient, we reads

$$\begin{aligned} p^0; \mathbb{T}_\alpha(u_0(t)) - 1 &= 0, \quad u_0(0) = 0, \\ p^1; \mathbb{T}_\alpha(u_1(t, c_1)) - \mathbb{T}_\alpha(u_0(t)) + 1 - c_1 [\mathbb{T}_\alpha(u_0(t)) + u_0^2(t) - 1] &= 0, \quad u_1(0) = 0, \\ p^2; \mathbb{T}_\alpha(u_2(t, c_1, c_2)) - \mathbb{T}_\alpha(u_1(t, c_1)) - c_2 [\mathbb{T}_\alpha(u_0(t)) - u_0^2(t) - 1] &= 0, \\ -c_1 [\mathbb{T}_\alpha(u_1(t, c_1)) + 2u_0(t)u_1(t, c_1)] &= 0, \quad u_2(0) = 0, \\ &\quad \vdots \end{aligned} \quad (4.11)$$

Corresponding solution of system of equations (4.11), are

$$\begin{aligned} u_0(t) &= \frac{1}{\alpha}t^\alpha, \\ u_1(t, c_1) &= \frac{1}{3}c_1\left(\frac{1}{\alpha}t^\alpha\right)^3, \\ u_2(t, c_1, c_2) &= \frac{1}{3}\left(\frac{1}{\alpha}t^\alpha\right)^3(c_1^2 + c_1 + c_2) + \frac{2}{15}\left(\frac{1}{\alpha}t^\alpha\right)^5c_1^2, \\ &\vdots \end{aligned}$$

Two-terms approximation to the solution of Eq. (4.9), will be obtained as follows

$$\begin{aligned} u^2(t, c_1, c_2) &= \left(\frac{1}{\alpha}t^\alpha + (0.6666666667c_1 + 0.3333333333c_1^2 \right. & (4.12) \\ &\left. + 0.3333333333c_2)\left(\frac{1}{\alpha}t^\alpha\right)^3 + 0.1333333333c_1^2\left(\frac{1}{\alpha}t^\alpha\right)^5. \end{aligned}$$

Table 4.2, shows the optimal values of the convergence control constants c_1 , c_2 and c_3 in $u^2(t, c_1, c_2)$ given in Eq. (4.12) for different values of α which can be obtained using the procedure mentioned in (3.9) up to (3.11). In Figure 4.3, the exact and approximate solutions of conformable fractional Riccati equation for $\alpha = 0.4, 0.6, 0.8$, and 1.0 are plotted.

Example 4.4. Consider the following fractional Riccati differential equation with initial value

$$T_\alpha u(t) = 2u(t) - u^2(t) + 1 = 0, 0 \leq t \leq 1, 0 < \alpha \leq 1, u(0) = 0. \quad (4.13)$$

The exact solution of Eq. (4.13), is $u(t) = \frac{\exp\left(\frac{2\sqrt{2}}{\alpha}t^\alpha\right) - 1}{(\sqrt{2}+1) + (\sqrt{2}-1)\exp\left(\frac{2\sqrt{2}}{\alpha}t^\alpha\right)}$.

By the proposed conformable fractional OHAM approach, we gives

$$(1 - p) [T_\alpha(v(t, p)) - 1] - H(p) [T_\alpha(v(t, p)) - 2v(t, p) + v^2(t, p) - 1] = 0. \quad (4.14)$$

Substituting Eqs. (3.4) and (3.5) into Eq. (4.14) and setting to zero the coefficient p , we reads

$$\begin{aligned} p^0; T_\alpha(u_0(t)) - 1 &= 0, \quad u_0(0) = 0, \\ p^1; T_\alpha(u_1(t, c_1)) - T_\alpha(u_0(t)) + 1 - c_1 [T_\alpha(u_0(t)) + u_0^2(t) - 2u_0(t) - 1] &= 0, \quad u_1(0) = 0, \\ p^2; T_\alpha(u_2(t, c_1, c_2)) - T_\alpha(u_1(t, c_1)) - c_2 [T_\alpha(u_0(t)) + u_0^2(t) - 2u_0(t) - 1] & \\ - c_1 [T_\alpha(u_1(t, c_1)) - 2u_1(t, c_1) + 2u_0(t)u_1(t, c_1)] &= 0, \quad u_2(0) = 0, \\ &\vdots \end{aligned} \quad (4.15)$$

Matching solution of Eqs. (4.15) are

$$\begin{aligned} u_0(t) &= \frac{1}{\alpha}t^\alpha, \\ u_1(t, c_1) &= c_1 \left[\frac{1}{3}\left(\frac{1}{\alpha}t^\alpha\right)^3 - \left(\frac{1}{\alpha}t^\alpha\right)^2 \right], \\ u_2(t, c_1, c_2) &= - (c_1^2 + c_1 + c_2) \left(\frac{1}{\alpha}t^\alpha\right)^2 + \left(\frac{1}{3}c_2 - \frac{2}{3}c_1^2 + \frac{1}{3}c_1\right) \left(\frac{1}{\alpha}t^\alpha\right)^3 + \frac{2}{15}c_1^2\left(\frac{1}{\alpha}t^\alpha\right)^5, \\ &\vdots \end{aligned}$$

Consequently, two-terms approximation to the solution of Eq. (4.13), will be obtained as the following

$$\begin{aligned} u^2(t, c_1, c_2) = & \left(\frac{1}{\alpha}t^\alpha\right) - [2c_1 + 0.9999999999c_1^2 + 0.9999999999c_2] \left(\frac{1}{\alpha}t^\alpha\right)^2 \\ & + [0.6666666666c_1 + 0.9999999999c_1^2 + 0.3333333333c_2] \left(\frac{1}{\alpha}t^\alpha\right)^3 \\ & - 0.6666666666c_1^2 \left(\frac{1}{\alpha}t^\alpha\right)^4 + 0.1333333333c_1^2 \left(\frac{1}{\alpha}t^\alpha\right)^5. \end{aligned} \quad (4.16)$$

Table 4.3, shows the optimal values of the convergence control constants c_1 , c_2 and c_3 in $u^2(t, c_1, c_2)$ given in Eq. (4.16) for different values of α which can be obtained using the procedure mentioned in (3.9) up to (3.11). In Figure 4.4, the exact solution and solution of conformable fractional OHAM of fractional Riccati equation, for $\alpha = 0.4, 0.6, 0.8,$ and 1.0 are plotted.

5. CONCLUSION

In this paper, optimal homotopy asymptotic method is applied to obtain an approximate solution of fractional differential equations. Conformable fractional derivatives are used for fractional derivative in this study. In comparison, of the results of applying conformable fractional derivatives with the results reported in (see [21, 22]), one learns that, CFD is a simple tool to find an approximate solution to a linear and a nonlinear fractional differential equation. What can one learn from the plots: The closer values of α , the larger convergence interval. To show the effectiveness of the method, fractional relaxation-oscillation equation, and Riccati fractional differential equation have been solved by conformable fractional optimal homotopy asymptotic method.

REFERENCES

- [1] R. Khalil, M.A. Horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, Journal of Computational and Applied Mathematics, Vol.264(2014), pp.65-70.
- [2] T. Abdeljawad, On conformable fractional calculus, Journal of Computational and Applied Mathematics, Vol.279(2015), pp.57-66.
- [3] F.B.M. Duarte, J.A.T. Machado, Chaotic phenomena and fractional- order dynamics in the trajectory control of redundant manipulators, Nonlinear Dynamics, Vol.29(2002), pp.342-362.
- [4] O.P. Agrawal, A general formulation and solution scheme for fractional optimal control problems, Nonlinear Dynamics, Vol.34(2004), pp.323-337.
- [5] N. Engheta, On fractional calculus and fractional multipoles in electromagnetism, IEEE Transactions on Antennas and Propagation, Vol.44(1996), pp.554-566.
- [6] R.L. Magin, Fractional calculus models of complex dynamics in biological tissues, Computers and Mathematics with Applications, Vol.59(2010), pp.1586-1593.
- [7] V.V. Kulish, J.L. Larg, Application of fractional calculus to fluid mechanics, Journal of Fluids Engineering, Vol.134(2002), doi:10.1115/1.1478062.
- [8] K.B. Oldhom, Fractional differential equations in electrochemistry, Advances Engineering Software, Vol.41(2010), pp.9-12.
- [9] V. Gaflychuk, B. Datsko, V. Meleshko, Mathematical modeling of time fractional reaction diffusion systems, Journal of Computational and Applied Mathematics, Vol.220(2008), pp.215-225.
- [10] A.R. Seadawy, Stability analysis solutions for nonlinear three-dimensional modified Korteweg-de Vries-Zakharov-Kuznetsov equation in a magnetized electron-positron plasma, Physica A: Statistical Mechanics and its Applications, Vol.455(2016), pp.44-51.
- [11] F.C. Meral, T.J. Royston, R. Magin, Fractional calculus in viscoelasticity:an experimental study, Communications in Nonlinear Science and Numerical Simulation, Vol.15(2010), pp.939-945.

- [12] F. Mainardi, Fractional calculus: some basic problems in continuum and statistical mechanics, in: A. Carpinteri, F. Mainardi (Eds.), *Fractals and Fractional calculus in continuum Mechanics*. New York: Springer-Verlag; 1997.
- [13] A.R. Seadawy, Nonlinear wave solutions of the three-dimensional Zakharov-Kuznetsov-Burgers equation in dusty plasma, *Physica A: Statistical Mechanics and its Applications*, Vol.439(2015), pp.1241-131.
- [14] V. DaftardarGejji, H. Jafari, Solving a multi-order fractional differential equation using Adomian Decomposition, *Applied Mathematics and Computation*, Vol.189(2007), pp.541-548.
- [15] B. Ghazanfari, A. Sepahvandzadeh, Adomian decomposition method for solving fractional Bratu-type equations, *Journal of Mathematics and Computer Science*, Vol.8(2014), pp.236-244.
- [16] O. Abdulaziz, I. Hashim, S. Momani, solving systems of fractional differential equations by homotopy perturbation method, *Physics Letters A*, Vol.372(2008), pp.451-459.
- [17] B. Ghazanfari, A.G. Ghazanfari, M. Fuladvand, Modification of the homotopy perturbation method for numerical solution of Nonlinear Wave and of Nonlinear Wave Equations., *Journal of Mathematics and Computer Science*, Vol.3(2011), pp.212-224.
- [18] M. Mahmoudi, M.V. Kazemi, Solving singular BVPs Ordinary Differential Equations by Modified Homotopy Perturbation Method, *Journal of Mathematics and Computer Science*, Vol.7(2013), pp.138-143.
- [19] M. Rabbani, New Homotopy Perturbation Method to Solve Non-Linear Problems, *Journal of Mathematics and Computer Science*, Vol.7(2013), pp.272-275.
- [20] I. Hashim, O. Abdulaziz, S. Momani, Homotopy Analysis Method for fractional IVPs, *Communications in Nonlinear Science and Numerical Simulation*, Vol.14(2009), pp.674-684.
- [21] M. Hamarsheh, A.I. Ismail, An Analytic Solution for Fractional Order Riccati Equations by Using Optimal Homotopy Asymptotic Method, *Applied Mathematical Sciences*, Vol.10(2016) No.23, pp.1131-1150.
- [22] M. Hamarsheh, A. Ismail, Z. Odibat, Optimal homotopy asymptotic method for solving fractional relaxation-oscillation equation, *Journal of Interpolation and Approximation in Scientific Computing*, Vol.2(2015), pp.98-111.
- [23] L.G. Yuan, Z. Alam, An optimal homotopy analysis method based on particle swarm optimization: application to fractional-order differential equation, *Journal of Applied Analysis and Computation*, Vol.6(2016) No.1, pp.103-118.
- [24] G. Wu, E.W.M. Lee, Fractional variational iteration method and its application, *Physics Letters A*, Vol.374(2010), pp.2506-2509.
- [25] Z. Odibat, S. Momani, V. S. Erturk, Generalized differential transform method: application to differential equations of fractional order, *Applied Mathematics and Computation*, Vol.197(2008), pp.467-477.
- [26] Y. Zhang, A finite difference method for fractional partial differential equation, *Applied Mathematics and Computation*, Vol.215(2009), pp.524-529.
- [27] H. Azizi, G.B. Loghmani, A numerical method for space fractional diffusion equations using a semi-discrete scheme and Chebyshev collocation method, *Journal of Mathematics and Computer Science*, Vol.8(2014), pp.2262-235.
- [28] A. Neamaty, B. Agheli, R. Darzi, Solving Fractional Partial Differential Equation by Using Wavelet Operational Method, *Journal of Mathematics and Computer Science*, Vol.7(2013), pp.230-240.
- [29] P. Rahimkhani, Y. Ordokhani, E. Babolian, Fractional-order Legendre wavelets and their applications for solving fractional-order differential equations with initial/boundary conditions, *Computational Methods for Differential Equations*, Vol.5(2017) No.2, pp.117-140.
- [30] A. Aghili, Solution to time fractional generalized KdV of order $2q+1$ and system of space fractional PDEs, *Computational Methods for Differential Equations*, Vol.5(2017) No.3, pp.246-255.
- [31] A. Neirameh, S. Shokooh, M. Eslami, Solutions structure of integrable families of Riccati equations and their applications to the perturbed nonlinear fractional Schrodinger equation, *Computational Methods for Differential Equations*, Vol.4(2016) No.4, pp.261-275.

- [32] M. Ekici, M. Mirzazadeh, M. Eslami, M. R. Belic, Optical soliton perturbation with fractional-temporal evolution by first integral method with conformable fractional derivatives, *Optik-International Journal for Light and Electron Optics*, Vol.127(2016) No.22, DOI: 10.1016/j.ijleo.2016.08.076.
- [33] M. Ilie, J. Biazar, Z. Ayati, The first integral method for solving some conformable fractional differential equations, *Optical and Quantum Electronics*, Vol.50(2018) No.2, <https://doi.org/10.1007/s11082-017-1307-x>.
- [34] M. Ilie, J. Biazar, Z. Ayati, General solution of Bernoulli and Riccati fractional differential equations based on conformable fractional derivative, *International Journal of Applied Mathematical Research*, Vol.6(2017) No.2, pp.49-51.
- [35] M. Ilie, J. Biazar, Z. Ayati, Application of the Lie Symmetry Analysis for second-order fractional differential equations, *Iranian Journal of Optimization*, Vol.9(2017) No.2, pp.79-83.
- [36] M. Ilie, J. Biazar, Z. Ayati, Analytical solutions for conformable fractional Bratu-type equations, *International Journal of Applied Mathematical Research*, Vol.7(2018) No.1, pp.15-19.

Figure 4.1. The exact and approximate solutions of fractional relaxation-oscillation (4.1) equation are plotted.




FIGURE 1. 2nd-order approximation of OHAM and Exact solution for Example 4.1.

Figures 4.2. The an exact and approximate solutions of fractional relaxation-oscillation equation for and are plotted.




FIGURE 2. 3rd-order approximation of OHAM and exact solution for Example 4.2.

Figures 4.3. The exact and approximate solutions of conformable fractional Riccati equation for and are plotted.




FIGURE 3. 2nd-order approximation of OHAM and exact solution for Example 4.3.

Figures 4.4. The exact solution and solution of conformable fractional OHAM of fractional Riccati equation, for and are plotted.




FIGURE 4. 2nd-order approximation of OHAM and exact solution for Example 4.4.

TABLE 4.1. Values of auxiliary parameters for the third-order OHAM solution of Eq. (4.5) for different orders.

Auxiliary parameters	$\alpha = 0.4$	$\alpha = 0.6$	$\alpha = 0.8$	$\alpha = 1.0$
c_1	-1.457903741	-1.208867087	-1.208867087	-1.155095209
c_2	0.1110279352	0.01085942753	0.01085942753	0.003733184279
c_3	-0.1200484251	0.001321301525	0.001321301525	0.001571139761

TABLE 4.2. Values of auxiliary parameters for the third-order OHAM solution of Eq. (4.9) for different orders.

Auxiliary parameters	$\alpha = 0.4$	$\alpha = 0.6$	$\alpha = 0.8$	$\alpha = 1.0$
c_1	-0.5315048975	0.6936479614	0.8425060981	-0.9175405542
c_2	0.0002051308039	-2.773945592	-3.369657214	0.0001113528473

TABLE 4.3. Values of auxiliary parameters for the third-order OHAM solution of Eq. (4.9) for different orders.

Auxiliary parameters	$\alpha = 0.4$	$\alpha = 0.6$	$\alpha = 0.8$	$\alpha = 1.0$
c_1	0.7431673291	-0.6503249572	-1.046574182	-1.0621151875
c_2	-3.154368233	-0.2366912237	-0.001518446383	0.002147348534

MOUSA ILIE

DEPARTMENT OF MATHEMATICS, GUILAN SCIENCE AND RESEARCH BRANCH, ISLAMIC AZAD UNIVERSITY, RASHT, IRAN

DEPARTMENT OF MATHEMATICS, RASHT BRANCH, ISLAMIC AZAD UNIVERSITY, RASHT, IRAN

E-mail address: ilie@iaurasht.ac.ir

JAFAR BIAZAR

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, UNIVERSITY OF GUILAN, P.O. BOX. 41335-1914, GUILAN, RASHT, IRAN

DEPARTMENT OF MATHEMATICS, RASHT BRANCH, ISLAMIC AZAD UNIVERSITY, RASHT, IRAN

E-mail address: biazar@iaurasht.ac.ir

ZAINAB AYATI

DEPARTMENT OF ENGINEERING SCIENCES, FACULTY OF TECHNOLOGY AND ENGINEERING EAST OF GUILAN, UNIVERSITY OF GUILAN, P.C. 44891-63157, RUDSAR-VAJARGAH, IRAN

E-mail address: ayati.zainab@gmail.com