

SOME NEW PARAMETERIZED INEQUALITIES FOR PREINVEX FUNCTIONS VIA GENERALIZED FRACTIONAL INTEGRAL OPERATORS AND THEIR APPLICATIONS

ARTION KASHURI, MUHAMMAD AMER LATIF

ABSTRACT. The authors have proved an identity with parameter for differentiable function with respect to another function via generalized integral operator. By applying the established identity, the generalized trapezium, Ostrowski and Simpson type integral inequalities have been discovered. Various special cases have been identified. Some applications of presented results to special means and new error estimates for the trapezium and midpoint quadrature formula have been analyzed. The ideas and techniques of this paper may stimulate further research in the field of integral inequalities.

1. INTRODUCTION

The following inequality, named Hermite–Hadamard inequality, is one of the most famous inequalities in the literature for convex functions.

Theorem 1.1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function and $a_1, a_2 \in I$ with $a_1 < a_2$. Then the following inequality holds:*

$$f\left(\frac{a_1 + a_2}{2}\right) \leq \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(x) dx \leq \frac{f(a_1) + f(a_2)}{2}. \quad (1.1)$$

This inequality (1.1) is also known as trapezium inequality.

The trapezium inequality has remained an area of great interest due to its wide applications in the field of mathematical analysis. Authors of recent decades have studied (1.1) in the premises of newly invented definitions due to motivation of convex function. Interested readers see the references [4]–[8],[17, 21, 22, 26, 27, 31, 33, 37],[39]–[42],[49, 51, 54, 55, 57, 58].

The following result is known in the literature as the Ostrowski inequality, see [34] and the references cited therein, which gives an upper bound for the approximation of the integral average $\frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(t) dt$ by the value $f(x)$ at point $x \in [a_1, a_2]$.

2010 *Mathematics Subject Classification.* 26A51, 26A33, 26D07, 26D10, 26D15.

Key words and phrases. Trapezium inequality, Ostrowski inequality, Simpson inequality, preinvexity, general fractional integrals.

Submitted January 6, 2020. Revised April 3, 2020.

Theorem 1.2. Let $f : I \rightarrow \mathbb{R}$ be a mapping differentiable on I° and let $a_1, a_2 \in I^\circ$ with $a_1 < a_2$. If $|f'(x)| \leq M$ for all $x \in [a, b]$, then

$$\left| f(x) - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(t) dt \right| \leq M(a_2 - a_1) \left[\frac{1}{4} + \frac{(x - \frac{a_1+a_2}{2})^2}{(a_2 - a_1)^2} \right], \quad \forall x \in [a_1, a_2]. \quad (1.2)$$

For other recent results concerning Ostrowski type inequalities, see [1]–[3],[9]–[16],[19, 23, 24],[34]–[36],[38],[43]–[45],[47, 48, 52, 56, 59]. Ostrowski inequality is playing a very important role in all the fields of mathematics, especially in the theory of approximations. Thus such inequalities were studied extensively by many researches and numerous generalizations, extensions and variants of them for various kind of functions like bounded variation, synchronous, Lipschitzian, monotonic, absolutely continuous and n -times differentiable mappings etc.

The following inequality is well known in the literature as Simpson's inequality.

Theorem 1.3. Let $f : [a_1, a_2] \rightarrow \mathbb{R}$ be four time differentiable on the interval (a_1, a_2) and having the fourth derivative bounded on (a_1, a_2) , that is $\|f^{(4)}\|_\infty = \sup_{x \in (a_1, a_2)} |f^{(4)}| < +\infty$. Then, we have

$$\left| \int_{a_1}^{a_2} f(x) dx - \frac{a_2 - a_1}{3} \left[\frac{f(a_1) + f(a_2)}{2} + 2f\left(\frac{a_1 + a_2}{2}\right) \right] \right| \leq \frac{1}{2880} \|f^{(4)}\|_\infty (a_2 - a_1)^5. \quad (1.3)$$

Inequality (1.3) gives an error bound for the classical Simpson quadrature formula, which is one of the most used quadrature formulae in practical applications. In recent years, various generalizations, extensions and variants of such inequalities have been obtained. For other recent results concerning Simpson type inequalities, see [32, 46, 53, 59].

In numerical analysis many quadrature rules have been established to approximate the definite integrals. Ostrowski inequality provides the bounds for many numerical quadrature rules, see [15, 16].

The aim of this paper is to establish trapezium, Ostrowski and Simpson type generalized integral inequalities for preinvex functions with respect to another function, some applications to special means and new error bounds for midpoint and trapezium quadrature formula. Interestingly, the special cases of presented results, are fractional integral inequalities. Therefore, it is important to summarize the study of fractional integrals.

At start, let us recall some mathematical preliminaries and definitions which will be helpful for further study.

Definition 1.4. [40] Let $f \in L[a_1, a_2]$. Then k -fractional integrals of order $\alpha, k > 0$ with $a_1 \geq 0$ are defined by

$$I_{a_1^+}^{\alpha, k} f(x) = \frac{1}{k\Gamma_k(\alpha)} \int_{a_1}^x (x-t)^{\frac{\alpha}{k}-1} f(t) dt, \quad x > a_1$$

and

$$I_{a_2}^{\alpha,k} f(x) = \frac{1}{k\Gamma_k(\alpha)} \int_x^{a_2} (t-x)^{\frac{\alpha}{k}-1} f(t) dt, \quad a_2 > x, \quad (1.4)$$

where $\Gamma_k(\cdot)$ is k -gamma function.

For $k = 1$, k -fractional integrals give Riemann-Liouville integrals. For $\alpha = k = 1$, k -fractional integrals give classical integrals.

Definition 1.5. [28, 29] Let $g : [a_1, a_2] \rightarrow \mathbb{R}$ be an increasing and positive monotone function on $[a_1, a_2]$, having a continuous derivative on (a_1, a_2) . The left-sided fractional integral of f with respect to g on $[a_1, a_2]$ of order $\alpha > 0$ is defined by:

$$I_{a_1+}^{\alpha,g} f(x) = \frac{1}{\Gamma(\alpha)} \int_{a_1}^x \frac{g'(u)f(u)}{[g(x) - g(u)]^{1-\alpha}} du, \quad x > a_1, \quad (1.5)$$

provided that the integral exists. The right-sided fractional integral of f with respect to g on $[a_1, a_2]$ of order $\alpha > 0$ is defined by:

$$I_{a_2-}^{\alpha,g} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^{a_2} \frac{g'(u)f(u)}{[g(u) - g(x)]^{1-\alpha}} du, \quad x < a_2, \quad (1.6)$$

provided that the integral exists.

Jleli and Samet in [21], proved the Hadamard type inequality for Riemann-Liouville fractional integral of a convex function f with respect to another function g .

Also in [49], Sarikaya and Ertuğral defined a function $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ satisfying the following conditions:

$$\int_0^1 \frac{\varphi(t)}{t} dt < +\infty, \quad (1.7)$$

$$\frac{1}{A} \leq \frac{\varphi(s)}{\varphi(r)} \leq A \text{ for } \frac{1}{2} \leq \frac{s}{r} \leq 2, \quad (1.8)$$

$$\frac{\varphi(r)}{r^2} \leq B \frac{\varphi(s)}{s^2} \text{ for } s \leq r, \quad (1.9)$$

$$\left| \frac{\varphi(r)}{r^2} - \frac{\varphi(s)}{s^2} \right| \leq C |r-s| \frac{\varphi(r)}{r^2} \text{ for } \frac{1}{2} \leq \frac{s}{r} \leq 2, \quad (1.10)$$

where $A, B, C > 0$ are independent of $r, s > 0$. If $\varphi(r)r^\alpha$ is increasing for some $\alpha \geq 0$ and $\frac{\varphi(r)}{r^\beta}$ is decreasing for some $\beta \geq 0$, then φ satisfies (1.7)-(1.10), see [50]. Therefore, the left-sided and right-sided generalized integral operators are defined as follows:

$${}_{a_1+} I_\varphi f(x) = \int_{a_1}^x \frac{\varphi(x-t)}{x-t} f(t) dt, \quad x > a_1, \quad (1.11)$$

$${}_{a_2-} I_\varphi f(x) = \int_x^{a_2} \frac{\varphi(t-x)}{t-x} f(t) dt, \quad x < a_2. \quad (1.12)$$

The most important feature of generalized integrals is that; they produce Riemann-Liouville fractional integrals, k -Riemann-Liouville fractional integrals, Katugampola fractional integrals, conformable fractional integrals, Hadamard fractional integrals, etc., see [20, 25, 49].

Recently, Farid in [18] generalised the above integral by introducing an increasing and positive monotone function g on $[a_1, a_2]$, having continuous derivative on (a_1, a_2) . The generalized fractional integral operator defined by Farid may be given as follows.

Definition 1.6. The left and right-sided generalized fractional integral of a function f with respect to another function g may be given as follows respectively:

$$G_{a_1+}^{\varphi,g} f(x) = \int_{a_1}^x \frac{\varphi(g(x) - g(u))}{g(x) - g(u)} g'(u) f(u) du, \quad x > a_1, \quad (1.13)$$

$$G_{a_2-}^{\varphi,g} f(x) = \int_x^{a_2} \frac{\varphi(g(u) - g(x))}{g(u) - g(x)} g'(u) f(u) du, \quad x < a_2. \quad (1.14)$$

This operator generalizes the various fractional integrals of a function f with respect to another function g .

The following special cases are focussed in our study.

(i) If we take $\varphi(u) = u$ then the operator (1.13) and (1.14) reduces to Riemann-Liouville integral of f with respect to function g .

$$I_{a_1+}^g f(x) = \int_{a_1}^x g'(u) f(u) du, \quad x > a_1, \quad (1.15)$$

$$I_{a_2-}^g f(x) = \int_x^{a_2} g'(u) f(u) du, \quad x < a_2. \quad (1.16)$$

If $g(u) = u$, then (1.15) and (1.16) will reduce to Riemann integral of f .

(ii) If we take $\varphi(u) = \frac{u^\alpha}{\Gamma(\alpha)}$ then the operator (1.13) and (1.14) reduces to Riemann-Liouville fractional integral of f with respect to function g .

$$I_{a_1+}^{\varphi,g} f(x) = \frac{1}{\Gamma(\alpha)} \int_{a_1}^x [g(x) - g(u)]^{\alpha-1} g'(u) f(u) du, \quad x > a_1, \quad (1.17)$$

$$I_{a_2-}^{\varphi,g} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^{a_2} [g(u) - g(x)]^{\alpha-1} g'(u) f(u) du, \quad x < a_2. \quad (1.18)$$

If $g(u) = u$, then (1.17) and (1.18) will reduce to left and right sided Riemann-Liouville fractional integrals of f respectively.

(iii) If we take $\varphi(u) = \frac{u^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ then the operator (1.13) and (1.14) reduces to k -Riemann-Liouville fractional integral of f with respect to function g .

$$I_{a_1+}^{\varphi,g,k} f(x) = \frac{1}{k\Gamma_k(\alpha)} \int_{a_1}^x [g(x) - g(u)]^{\frac{\alpha}{k}-1} g'(u) f(u) du, \quad x > a_1, \quad (1.19)$$

$$I_{a_2-}^{\varphi,g,k} f(x) = \frac{1}{k\Gamma_k(\alpha)} \int_x^{a_2} [g(u) - g(x)]^{\frac{\alpha}{k}-1} g'(u) f(u) du, \quad x < a_2. \quad (1.20)$$

If $g(u) = u$, then these operators in (1.19) and (1.20) reduces to k -fractional integral operators given in [40].

(iv) If we take $\varphi_g(u) = u(g(a_2) - u)^{\alpha-1}$ for $\alpha \in (0, 1)$, then the operator given in (1.13) and (1.14) reduces to conformable fractional integral operator of f with respect to a function g .

$$K_{a_1}^{\alpha, g} f(x) = \int_{a_1}^x [g(u)]^{\alpha-1} g'(u) f(u) du, \quad x > a_1. \quad (1.21)$$

This operator (1.21) generalizes conformable fractional integral operator which was given by Khalil et al. in [30].

(v) If we take $\varphi(u) = \frac{u}{\alpha} \exp(-Au)$, where $A = \frac{1-\alpha}{\alpha}$ and $\alpha \in (0, 1)$, then the operator given in (1.13) and (1.14) reduces to fractional integral operator of f with respect to function g with exponential kernel.

$$J_{a_1+}^{\alpha, g} f(x) = \frac{1}{\alpha} \int_{a_1}^x \exp(-A(g(x) - g(u))) g'(u) f(u) du, \quad x > a_1, \quad (1.22)$$

$$J_{a_2-}^{\alpha, g} f(x) = \frac{1}{\alpha} \int_x^{a_2} \exp(-A(g(x) - g(u))) g'(u) f(u) du, \quad x < a_2. \quad (1.23)$$

Operators in (1.22) and (1.23) generalizes fractional integral operator with exponential kernel which was introduced by Kirane and Torebek in [31].

Motivated by the above literatures, the main objective of this paper is to discover in section 2, an interesting identity with parameter in order to study some new bounds regarding trapezium, Ostrowski and Simpson type integral inequalities. By using the established identity as an auxiliary result, some new estimates for trapezium, Ostrowski and Simpson type integral inequalities via generalized integrals are obtained. It is pointed out that some new fractional integral inequalities have been deduced from main results. In section 3, some applications to special means and new error estimates for the midpoint and trapezium quadrature formula are given. The ideas and techniques of this paper may stimulate further research in the field of integral inequalities.

2. MAIN RESULTS

Throughout this study, let $P = [ma_1, a_2]$, where $a_1 < a_2$ and $m \in (0, 1]$. Also, let $\eta : P \times P \rightarrow \mathbb{R}$ and $t \in [0, 1]$. For brevity, we define

$$\begin{aligned} \Lambda_m^{\varphi, g}(x, t) &= \int_0^t \frac{\varphi(g(ma_1 + u\eta(x, ma_1)) - g(ma_1))}{g(ma_1 + u\eta(x, ma_1)) - g(ma_1)} \\ &\quad \times g'(ma_1 + u\eta(x, ma_1)) du < +\infty \end{aligned} \quad (2.1)$$

and

$$\begin{aligned} \Delta_m^{\varphi, g}(x, t) &= \int_t^1 \frac{\varphi(g(mx + \eta(a_2, mx)) - g(mx + u\eta(a_2, mx)))}{g(mx + \eta(a_2, mx)) - g(mx + u\eta(a_2, mx))} \\ &\quad \times g'(mx + u\eta(a_2, mx)) du < +\infty, \end{aligned} \quad (2.2)$$

where g is an increasing and positive monotone function on P , having continuous derivative on $P^\circ = (ma_1, a_2)$.

For establishing some new results regarding general fractional integrals we need to prove the following lemma.

Lemma 2.1. Let $f : P \rightarrow \mathbb{R}$ be a differentiable mapping on P° and $\lambda \in \mathbb{R}$. If $f' \in L(P)$ and $\eta(a_2, ma_1) > 0$, then the following identity for generalized fractional integrals hold:

$$\begin{aligned}
& \frac{\eta(x, ma_1)f(ma_1 + t\eta(x, ma_1)) + \eta(a_2, mx)f(mx)}{\eta(a_2, ma_1)} \\
& - \frac{\lambda}{\eta(a_2, ma_1)} \times \left[\frac{\eta(x, ma_1)f(ma_1 + t\eta(x, ma_1))}{\Lambda_m^{\varphi, g}(x, 1)} + \frac{\eta(a_2, mx)f(mx)}{\Delta_m^{\varphi, g}(x, 0)} \right] \\
& + \frac{\lambda}{\eta(a_2, ma_1)} \times \left[\frac{\eta(x, ma_1)f(ma_1)}{\Lambda_m^{\varphi, g}(x, 1)} + \frac{\eta(a_2, mx)f(mx + \eta(a_2, mx))}{\Delta_m^{\varphi, g}(x, 0)} \right] \\
& - \frac{1}{\eta(a_2, ma_1)} \times \left[\frac{G_{(ma_1 + \eta(x, ma_1))}^{\varphi, g}f(ma_1)}{\Lambda_m^{\varphi, g}(x, 1)} + \frac{G_{(mx)}^{\varphi, g}f(mx + \eta(a_2, mx))}{\Delta_m^{\varphi, g}(x, 0)} \right] \\
& = \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \times \int_0^1 [\Lambda_m^{\varphi, g}(x, t) - \lambda] f'(ma_1 + t\eta(x, ma_1)) dt \quad (2.3) \\
& - \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \times \int_0^1 [\Delta_m^{\varphi, g}(x, t) - \lambda] f'(mx + t\eta(a_2, mx)) dt.
\end{aligned}$$

We denote

$$\begin{aligned}
T_{f, \Lambda_m^{\varphi, g}, \Delta_m^{\varphi, g}}(\lambda; x, a_1, a_2) &= \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \\
&\times \int_0^1 [\Lambda_m^{\varphi, g}(x, t) - \lambda] f'(ma_1 + t\eta(x, ma_1)) dt \\
&- \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \times \int_0^1 [\Delta_m^{\varphi, g}(x, t) - \lambda] f'(mx + t\eta(a_2, mx)) dt.
\end{aligned} \quad (2.4)$$

Proof. Integrating by parts (2.4) and changing the variables of integration, we have

$$\begin{aligned}
T_{f, \Lambda_m^{\varphi, g}, \Delta_m^{\varphi, g}}(\lambda; x, a_1, a_2) &= \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \\
&\times \left\{ \int_0^1 \Lambda_m^{\varphi, g}(x, t) f'(ma_1 + t\eta(x, ma_1)) dt - \lambda \int_0^1 f'(ma_1 + t\eta(x, ma_1)) dt \right\} \\
&- \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \\
&\times \left\{ \int_0^1 \Delta_m^{\varphi, g}(x, t) f'(mx + t\eta(a_2, mx)) dt - \lambda \int_0^1 f'(mx + t\eta(a_2, mx)) dt \right\} \\
&= \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \times \left\{ \frac{\Lambda_m^{\varphi, g}(x, t)f(ma_1 + t\eta(x, ma_1))}{\eta(x, ma_1)} \Big|_0^1 - \frac{1}{\eta(x, ma_1)} \right. \\
&\times \int_0^1 \frac{\varphi(g(ma_1 + t\eta(x, ma_1)) - g(ma_1))}{g(ma_1 + t\eta(x, ma_1)) - g(ma_1)} g'(ma_1 + t\eta(x, ma_1)) f(ma_1 + t\eta(x, ma_1)) dt \\
&\left. - \frac{\lambda}{\eta(x, ma_1)} f(ma_1 + t\eta(x, ma_1)) \Big|_0^1 \right\} - \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \\
&\times \left\{ \frac{\Delta_m^{\varphi, g}(x, t)f(mx + t\eta(a_2, mx))}{\eta(a_2, mx)} \Big|_0^1 \right\}
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\eta(a_2, mx)} \times \int_0^1 \frac{\varphi(g(mx + \eta(a_2, mx)) - g(mx + t\eta(a_2, mx)))}{g(mx + \eta(a_2, mx)) - g(mx + t\eta(a_2, mx))} \\
& \times g'(mx + t\eta(a_2, mx)) f(mx + t\eta(a_2, mx)) dt - \frac{\lambda}{\eta(a_2, mx)} f(mx + t\eta(a_2, mx)) \Big|_0^1 \Big\} \\
& = \frac{\eta(x, ma_1) f(mx_1 + t\eta(x, ma_1)) + \eta(a_2, mx) f(mx)}{\eta(a_2, ma_1)} \\
& - \frac{\lambda}{\eta(a_2, ma_1)} \times \left[\frac{\eta(x, ma_1) f(mx_1 + t\eta(x, ma_1))}{\Lambda_m^{\varphi,g}(x, 1)} + \frac{\eta(a_2, mx) f(mx)}{\Delta_m^{\varphi,g}(x, 0)} \right] \\
& + \frac{\lambda}{\eta(a_2, ma_1)} \times \left[\frac{\eta(x, ma_1) f(mx_1)}{\Lambda_m^{\varphi,g}(x, 1)} + \frac{\eta(a_2, mx) f(mx + \eta(a_2, mx))}{\Delta_m^{\varphi,g}(x, 0)} \right] \\
& - \frac{1}{\eta(a_2, ma_1)} \times \left[\frac{G_{(ma_1 + \eta(x, ma_1))}^{\varphi,g} f(mx_1)}{\Lambda_m^{\varphi,g}(x, 1)} + \frac{G_{(mx)}^{\varphi,g} f(mx + \eta(a_2, mx))}{\Delta_m^{\varphi,g}(x, 0)} \right].
\end{aligned}$$

This completes the proof of our Lemma 2.1. \square

Remark 2.2. **a:** Taking $m = 1, \lambda = 0, \eta(x, ma_1) = x - ma_1, \eta(a_2, mx) = a_2 - mx, \eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Lemma 2.1, we have the following Ostrowski type identity:

$$T_f(x, a_1, a_2) = f(x) - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(t) dt.$$

b: Taking $m = 1, \lambda = 1, \eta(x, ma_1) = x - ma_1, \eta(a_2, mx) = a_2 - mx, \eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Lemma 2.1, we get the following Hermite-Hadamard type identity:

$$\overline{T}_f(x, a_1, a_2) = \frac{(x - a_1)f(a_1) + (a_2 - x)f(a_2)}{a_2 - a_1} - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(t) dt.$$

c: Taking $m = 1, x = \frac{a_1 + a_2}{2}, \eta(x, ma_1) = x - ma_1, \eta(a_2, mx) = a_2 - mx, \eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Lemma 2.1, we obtain the following Simpson type identity:

$$T_f(\lambda; a_1, a_2) = \lambda \left[\frac{f(a_1) + f(a_2)}{2} \right] + (1 - \lambda) f\left(\frac{a_1 + a_2}{2}\right) - \frac{1}{a_2 - a_1} \int_{a_1}^{a_2} f(t) dt.$$

Theorem 2.3. Let $f : P \rightarrow \mathbb{R}$ be a differentiable mapping on P° and $\lambda \in [0, 1]$. If $|f'|^q$ is preinvex on P and $\eta(a_2, ma_1) > 0$, then for $q > 1$ and $p^{-1} + q^{-1} = 1$, the following inequality for generalized fractional integrals hold:

$$\begin{aligned}
& |T_{f, \Lambda_m^{\varphi,g}, \Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| \\
& \leq \frac{\eta^2(x, ma_1)}{\sqrt[q]{2} \Lambda_m^{\varphi,g}(x, 1) \eta(a_2, ma_1)} \sqrt[p]{B_{\Lambda_m^{\varphi,g}}^{\varphi,g}(x; \lambda, p)} \times \sqrt[q]{|f'(ma_1)|^q + |f'(x)|^q} \quad (2.5) \\
& + \frac{\eta^2(a_2, mx)}{\sqrt[q]{2} \Delta_m^{\varphi,g}(x, 0) \eta(a_2, ma_1)} \sqrt[p]{B_{\Delta_m^{\varphi,g}}^{\varphi,g}(x; \lambda, p)} \times \sqrt[q]{|f'(mx)|^q + |f'(a_2)|^q},
\end{aligned}$$

where

$$B_{\Lambda_m^{\varphi,g}}^{\varphi,g}(x; \lambda, p) = \int_0^1 \left| \Lambda_m^{\varphi,g}(x, t) - \lambda \right|^p dt, \quad B_{\Delta_m^{\varphi,g}}^{\varphi,g}(x; \lambda, p) = \int_0^1 \left| \Delta_m^{\varphi,g}(x, t) - \lambda \right|^p dt. \quad (2.6)$$

Proof. From Lemma 2.1, preinvexity of $|f'|^q$, Hölder inequality and properties of the modulus, we have

$$\begin{aligned}
|T_{f,\Lambda_m^{\varphi,g},\Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| &\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \\
&\quad \times \int_0^1 |\Lambda_m^{\varphi,g}(x, t) - \lambda| |f'(ma_1 + t\eta(x, ma_1))| dt \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \times \int_0^1 |\Delta_m^{\varphi,g}(x, t) - \lambda| |f'(mx + t\eta(a_2, mx))| dt \\
&\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \times \left(\int_0^1 |\Lambda_m^{\varphi,g}(x, t) - \lambda|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(ma_1 + t\eta(x, ma_1))|^q dt \right)^{\frac{1}{q}} \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \times \left(\int_0^1 |\Delta_m^{\varphi,g}(x, t) - \lambda|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |f'(mx + t\eta(a_2, mx))|^q dt \right)^{\frac{1}{q}} \\
&\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \sqrt[p]{B_{\Lambda_m}^{\varphi,g}(x; \lambda, p)} \left(\int_0^1 [(1-t)|f'(ma_1)|^q + t|f'(x)|^q] dt \right)^{\frac{1}{q}} \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \sqrt[p]{B_{\Delta_m}^{\varphi,g}(x; \lambda, p)} \left(\int_0^1 [(1-t)|f'(mx)|^q + t|f'(a_2)|^q] dt \right)^{\frac{1}{q}} \\
&= \frac{\eta^2(x, ma_1)}{\sqrt[3]{2}\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \sqrt[p]{B_{\Lambda_m}^{\varphi,g}(x; \lambda, p)} \times \sqrt[3]{|f'(ma_1)|^q + |f'(x)|^q} \\
&\quad + \frac{\eta^2(a_2, mx)}{\sqrt[3]{2}\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \sqrt[p]{B_{\Delta_m}^{\varphi,g}(x; \lambda, p)} \times \sqrt[3]{|f'(mx)|^q + |f'(a_2)|^q}.
\end{aligned}$$

The proof of Theorem 2.3 is completed. \square

We point out some special cases of Theorem 2.3 as follows:

Corollary 2.4. *Taking $p = q = 2$ in Theorem 2.3, we have*

$$\begin{aligned}
&|T_{f,\Lambda_m^{\varphi,g},\Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| \\
&\leq \frac{\eta^2(x, ma_1)}{\sqrt{2}\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \sqrt{B_{\Lambda_m}^{\varphi,g}(x; \lambda, 2)} \times \sqrt{|f'(ma_1)|^2 + |f'(x)|^2} \quad (2.7) \\
&\quad + \frac{\eta^2(a_2, mx)}{\sqrt{2}\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \sqrt{B_{\Delta_m}^{\varphi,g}(x; \lambda, 2)} \times \sqrt{|f'(mx)|^2 + |f'(a_2)|^2}.
\end{aligned}$$

Corollary 2.5. *Taking $|f'| \leq K$ in Theorem 2.3, we get*

$$\begin{aligned}
&|T_{f,\Lambda_m^{\varphi,g},\Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| \leq \frac{K}{\eta(a_2, ma_1)} \\
&\quad \times \left[\frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)} \sqrt[p]{B_{\Lambda_m}^{\varphi,g}(x; \lambda, p)} + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)} \sqrt[p]{B_{\Delta_m}^{\varphi,g}(x; \lambda, p)} \right]. \quad (2.8)
\end{aligned}$$

Corollary 2.6. Taking $m = 1$, $\lambda = 0$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.3, we obtain the following Ostrowski type inequality:

$$\begin{aligned} |T_f(x, a_1, a_2)| &\leq \frac{1}{\sqrt[4]{2}\sqrt[4]{p+1}(a_2 - a_1)} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{|f'(a_1)|^q + |f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{|f'(x)|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.9)$$

Corollary 2.7. Taking $x = \frac{a_1+a_2}{2}$ in Corollary 2.6, we have the following midpoint type inequality:

$$\begin{aligned} |T_f(a_1, a_2)| &\leq \frac{(a_2 - a_1)}{4\sqrt[4]{2}\sqrt[4]{p+1}} \\ &\times \left\{ \sqrt[4]{|f'(a_1)|^q + \left| f' \left(\frac{a_1 + a_2}{2} \right) \right|^q} + \sqrt[4]{\left| f' \left(\frac{a_1 + a_2}{2} \right) \right|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.10)$$

Corollary 2.8. Taking $m = 1$, $\lambda = 1$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.3, we get the following trapezium type inequality:

$$\begin{aligned} |\overline{T}_f(x, a_1, a_2)| &\leq \frac{1}{\sqrt[4]{2}\sqrt[4]{p+1}(a_2 - a_1)} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{|f'(a_1)|^q + |f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{|f'(x)|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.11)$$

Corollary 2.9. Taking $m = 1$, $\lambda = \frac{1}{3}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.3, we obtain the following Simpson type inequality:

$$\begin{aligned} \left| T_f \left(\frac{1}{3}; a_1, a_2 \right) \right| &\leq \frac{1}{\sqrt[4]{2}(a_2 - a_1)} \sqrt[p]{\frac{2^{p+1} + 1}{3^{p+1}(p+1)}} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{|f'(a_1)|^q + |f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{|f'(x)|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.12)$$

Corollary 2.10. Taking $\lambda = 0$ and $\varphi(t) = t$ in Theorem 2.3, we have

$$\begin{aligned} |T_{f, \Delta_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\sqrt[4]{2}\eta(a_2, ma_1)} \\ &\times \left\{ \sqrt[4]{\eta(x, ma_1)} \sqrt[p]{B_1^g(x; p)} \times \sqrt[4]{|f'(ma_1)|^q + |f'(x)|^q} \right. \\ &\left. + \sqrt[4]{\eta(a_2, mx)} \sqrt[p]{B_2^g(x; p)} \times \sqrt[4]{|f'(mx)|^q + |f'(a_2)|^q} \right\}, \end{aligned} \quad (2.13)$$

where

$$B_1^g(x; p) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} [g(t) - g(ma_1)]^p dt \quad (2.14)$$

and

$$B_2^g(x; p) = \int_{mx}^{mx + \eta(a_2, mx)} [g(mx + \eta(a_2, mx)) - g(t)]^p dt. \quad (2.15)$$

Corollary 2.11. Taking $\lambda = 0$ and $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$ in Theorem 2.3, we get

$$\begin{aligned} |T_{f,\Lambda_m^g,\Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\sqrt[2]{2}\eta(a_2, ma_1)} \\ &\times \left\{ \sqrt[q]{\eta(x, ma_1)} \sqrt[p]{B_3^g(x; p, \alpha)} \times \sqrt[q]{|f'(ma_1)|^q + |f'(x)|^q} \right. \\ &\left. + \sqrt[q]{\eta(a_2, mx)} \sqrt[p]{B_4^g(x; p, \alpha)} \times \sqrt[q]{|f'(mx)|^q + |f'(a_2)|^q} \right\}, \end{aligned} \quad (2.16)$$

where

$$B_3^g(x; p, \alpha) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} [g(t) - g(ma_1)]^{p\alpha} dt \quad (2.17)$$

and

$$B_4^g(x; p, \alpha) = \int_{mx}^{mx + \eta(a_2, mx)} [g(mx + \eta(a_2, mx)) - g(t)]^{p\alpha} dt. \quad (2.18)$$

Corollary 2.12. Taking $\lambda = 0$ and $\varphi(t) = \frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ in Theorem 2.3, we obtain

$$|T_{f,\Lambda_m^g,\Delta_m^g}(0; x, a_1, a_2)| \leq \frac{1}{\sqrt[2]{2}\eta(a_2, ma_1)} \quad (2.19)$$

$$\begin{aligned} &\times \left\{ \sqrt[q]{\eta(x, ma_1)} \sqrt[p]{B_5^g(x; p, \alpha, k)} \times \sqrt[q]{|f'(ma_1)|^q + |f'(x)|^q} \right. \\ &\left. + \sqrt[q]{\eta(a_2, mx)} \sqrt[p]{B_6^g(x; p, \alpha, k)} \times \sqrt[q]{|f'(mx)|^q + |f'(a_2)|^q} \right\}, \end{aligned}$$

where

$$B_5^g(x; p, \alpha, k) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} [g(t) - g(ma_1)]^{\frac{p\alpha}{k}} dt \quad (2.20)$$

and

$$B_6^g(x; p, \alpha, k) = \int_{mx}^{mx + \eta(a_2, mx)} [g(mx + \eta(a_2, mx)) - g(t)]^{\frac{p\alpha}{k}} dt. \quad (2.21)$$

Corollary 2.13. Taking $\lambda = 0$, $\forall u \in [0, t]$, $\varphi_g(x, t) = t(g(ma_1 + \eta(x, ma_1)) - t)^{\alpha-1}$ and $\forall u \in [t, 1]$, $\varphi_g(x, t) = t(g(mx + \eta(a_2, mx)) - t)^{\alpha-1}$ in Theorem 2.3, we have

$$\begin{aligned} |T_{f,\Lambda_m^g,\Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{\eta^{\frac{q+1}{q}}(x, ma_1)}{\sqrt[2]{2}[g(ma_1 + \eta(x, ma_1)) - g(ma_1)]\eta(a_2, ma_1)} \quad (2.22) \\ &\times \sqrt[p]{B_7^g(x; p)} \times \sqrt[q]{|f'(ma_1)|^q + |f'(x)|^q} \\ &+ \frac{\eta^{\frac{q+1}{q}}(a_2, mx)}{\sqrt[2]{2}[g^\alpha(mx + \eta(a_2, mx)) - g^\alpha(mx)]\eta(a_2, ma_1)} \\ &\times \sqrt[p]{B_8^g(x; p, \alpha)} \times \sqrt[q]{|f'(mx)|^q + |f'(a_2)|^q}, \end{aligned}$$

where

$$B_7^g(x; p) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} [g(t) - g(ma_1)]^p dt \quad (2.23)$$

and

$$B_8^g(x; p, \alpha) = \int_{mx}^{mx+\eta(a_2, mx)} \left[g^\alpha(mx + \eta(a_2, mx)) - g^\alpha(t) \right]^p dt. \quad (2.24)$$

Corollary 2.14. *Taking $\lambda = 0$ and $\varphi(t) = \frac{t}{\alpha} \exp(-At)$, where $A = \frac{1-\alpha}{\alpha}$, in Theorem 2.3, we get*

$$\begin{aligned} |T_{f, \Lambda_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{\eta^{\frac{q+1}{q}}(x, ma_1)}{\sqrt[q]{2\eta(a_2, ma_1)}} \sqrt[p]{B_9^g(x; p, A)} \\ &\quad \times \sqrt[q]{|f'(ma_1)|^q + |f'(x)|^q} \\ &+ \frac{\eta^{\frac{q+1}{q}}(a_2, mx)}{\sqrt[q]{2\eta(a_2, ma_1)}} \sqrt[p]{B_{10}^g(x; p, A)} \times \sqrt[q]{|f'(mx)|^q + |f'(a_2)|^q}, \end{aligned} \quad (2.25)$$

where

$$B_9^g(x; p, A) = \int_{ma_1}^{ma_1+\eta(x, ma_1)} \left\{ 1 - \exp [A(g(ma_1) - g(t))] \right\}^p dt \quad (2.26)$$

and

$$B_{10}^g(x; p, A) = \int_{mx}^{mx+\eta(a_2, mx)} \left\{ 1 - \exp [A(g(t) - g(mx + \eta(a_2, mx)))] \right\}^p dt. \quad (2.27)$$

Theorem 2.15. *Let $f : P \rightarrow \mathbb{R}$ be a differentiable mapping on P° and $\lambda \in [0, 1]$. If $|f'|^q$ is preinvex on P and $\eta(a_2, ma_1) > 0$, then for $q \geq 1$, the following inequality for generalized fractional integrals hold:*

$$\begin{aligned} |T_{f, \Lambda_m^{\varphi, g}, \Delta_m^{\varphi, g}}(\lambda; x, a_1, a_2)| &\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \left[B_{\Lambda_m^g}^{\varphi, g}(x; \lambda, 1) \right]^{1-\frac{1}{q}} \\ &\quad \times \sqrt[q]{\left[B_{\Lambda_m^g}^{\varphi, g}(x; \lambda, 1) - E_{\Lambda_m^g}^{\varphi, g}(x; \lambda) \right] |f'(ma_1)|^q + E_{\Lambda_m^g}^{\varphi, g}(x; \lambda) |f'(x)|^q} \\ &\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \left[B_{\Delta_m^g}^{\varphi, g}(x; \lambda, 1) \right]^{1-\frac{1}{q}} \\ &\quad \times \sqrt[q]{\left[B_{\Delta_m^g}^{\varphi, g}(x; \lambda, 1) - G_{\Delta_m^g}^{\varphi, g}(x; \lambda) \right] |f'(mx)|^q + G_{\Delta_m^g}^{\varphi, g}(x; \lambda) |f'(a_2)|^q}, \end{aligned} \quad (2.28)$$

where

$$E_{\Lambda_m^g}^{\varphi, g}(x; \lambda) = \int_0^1 t \left| \Lambda_m^{\varphi, g}(x, t) - \lambda \right| dt, \quad G_{\Delta_m^g}^{\varphi, g}(x; \lambda) = \int_0^1 t \left| \Delta_m^{\varphi, g}(x, t) - \lambda \right| dt \quad (2.29)$$

and $B_{\Lambda_m^g}^{\varphi, g}(x; \lambda, 1)$, $B_{\Delta_m^g}^{\varphi, g}(x; \lambda, 1)$ are defined as in Theorem 2.3.

Proof. From Lemma 2.1, preinvexity of $|f'|^q$, the well-known power mean inequality and properties of the modulus, we have

$$\begin{aligned} |T_{f, \Lambda_m^{\varphi, g}, \Delta_m^{\varphi, g}}(\lambda; x, a_1, a_2)| &\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi, g}(x, 1)\eta(a_2, ma_1)} \\ &\quad \times \int_0^1 \left| \Lambda_m^{\varphi, g}(x, t) - \lambda \right| \left| f'(ma_1 + t\eta(x, ma_1)) \right| dt \\ &+ \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi, g}(x, 0)\eta(a_2, ma_1)} \times \int_0^1 \left| \Delta_m^{\varphi, g}(x, t) - \lambda \right| \left| f'(mx + t\eta(a_2, mx)) \right| dt \end{aligned}$$

$$\begin{aligned}
&\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \\
&\times \left(\int_0^1 |\Lambda_m^{\varphi,g}(x, t) - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |\Lambda_m^{\varphi,g}(x, t) - \lambda| |f'(ma_1 + t\eta(x, ma_1))|^q dt \right)^{\frac{1}{q}} \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \\
&\times \left(\int_0^1 |\Delta_m^{\varphi,g}(x, t) - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |\Delta_m^{\varphi,g}(x, t) - \lambda| |f'(mx + t\eta(a_2, mx))|^q dt \right)^{\frac{1}{q}} \\
&\leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \sqrt[q]{B_{\Lambda_m}^{\varphi,g}(x; \lambda, p)} \\
&\times \left(\int_0^1 |\Lambda_m^{\varphi,g}(x, t) - \lambda| [(1-t)|f'(ma_1)|^q + t|f'(x)|^q] dt \right)^{\frac{1}{q}} \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \sqrt[q]{B_{\Delta_m}^{\varphi,g}(x; \lambda, p)} \\
&\times \left(\int_0^1 |\Delta_m^{\varphi,g}(x, t) - \lambda| [(1-t)|f'(mx)|^q + t|f'(a_2)|^q] dt \right)^{\frac{1}{q}} \\
&= \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \left[B_{\Lambda_m}^{\varphi,g}(x; \lambda, 1) \right]^{1-\frac{1}{q}} \\
&\times \sqrt[q]{ \left[B_{\Lambda_m}^{\varphi,g}(x; \lambda, 1) - E_{\Lambda_m}^{\varphi,g}(x; \lambda) \right] |f'(ma_1)|^q + E_{\Lambda_m}^{\varphi,g}(x; \lambda) |f'(x)|^q } \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \left[B_{\Delta_m}^{\varphi,g}(x; \lambda, 1) \right]^{1-\frac{1}{q}} \\
&\times \sqrt[q]{ \left[B_{\Delta_m}^{\varphi,g}(x; \lambda, 1) - G_{\Delta_m}^{\varphi,g}(x; \lambda) \right] |f'(mx)|^q + G_{\Delta_m}^{\varphi,g}(x; \lambda) |f'(a_2)|^q }.
\end{aligned}$$

The proof of Theorem 2.15 is completed. \square

We point out some special cases of Theorem 2.15 as follows:

Corollary 2.16. *Taking $q = 1$ in Theorem 2.15, we have*

$$\begin{aligned}
&|T_{f, \Lambda_m^{\varphi,g}, \Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| \leq \frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)\eta(a_2, ma_1)} \tag{2.30} \\
&\times \left[(B_{\Lambda_m}^{\varphi,g}(x; \lambda, 1) - E_{\Lambda_m}^{\varphi,g}(x; \lambda)) |f'(ma_1)| + E_{\Lambda_m}^{\varphi,g}(x; \lambda) |f'(x)| \right] \\
&\quad + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)\eta(a_2, ma_1)} \\
&\times \left[(B_{\Delta_m}^{\varphi,g}(x; \lambda, 1) - G_{\Delta_m}^{\varphi,g}(x; \lambda)) |f'(mx)| + G_{\Delta_m}^{\varphi,g}(x; \lambda) |f'(a_2)| \right].
\end{aligned}$$

Corollary 2.17. *Taking $|f'| \leq K$ in Theorem 2.15, we get*

$$\begin{aligned}
&|T_{f, \Lambda_m^{\varphi,g}, \Delta_m^{\varphi,g}}(\lambda; x, a_1, a_2)| \leq \frac{K}{\eta(a_2, ma_1)} \tag{2.31} \\
&\times \left[\frac{\eta^2(x, ma_1)}{\Lambda_m^{\varphi,g}(x, 1)} B_{\Lambda_m}^{\varphi,g}(x; \lambda, 1) + \frac{\eta^2(a_2, mx)}{\Delta_m^{\varphi,g}(x, 0)} B_{\Delta_m}^{\varphi,g}(x; \lambda, 1) \right].
\end{aligned}$$

Corollary 2.18. Taking $m = 1$, $\lambda = 0$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.15, we obtain the following Ostrowski type inequality:

$$\begin{aligned} |T_f(x, a_1, a_2)| &\leq \frac{1}{2\sqrt[4]{3}(a_2 - a_1)} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{|f'(a_1)|^q + 2|f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{2|f'(x)|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.32)$$

Corollary 2.19. Taking $x = \frac{a_1 + a_2}{2}$ in Corollary 2.18, we have the following midpoint type inequality:

$$\begin{aligned} |T_f(a_1, a_2)| &\leq \frac{(a_2 - a_1)}{8\sqrt[4]{3}} \\ &\times \left\{ \sqrt[4]{|f'(a_1)|^q + 2\left|f'\left(\frac{a_1 + a_2}{2}\right)\right|^q} + \sqrt[4]{2\left|f'\left(\frac{a_1 + a_2}{2}\right)\right|^q + |f'(a_2)|^q} \right\}. \end{aligned} \quad (2.33)$$

Corollary 2.20. Taking $m = 1$, $\lambda = 1$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.15, we get the following trapezium type inequality:

$$\begin{aligned} |\overline{T}_f(x, a_1, a_2)| &\leq \frac{1}{2\sqrt[4]{3}(a_2 - a_1)} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{2|f'(a_1)|^q + |f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{|f'(x)|^q + 2|f'(a_2)|^q} \right\}. \end{aligned} \quad (2.34)$$

Corollary 2.21. Taking $m = 1$, $\lambda = \frac{1}{3}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ in Theorem 2.15, we obtain the following Simpson type inequality:

$$\begin{aligned} \left| T_f \left(\frac{1}{3}; a_1, a_2 \right) \right| &\leq \frac{1}{2\sqrt[4]{243}(a_2 - a_1)} \\ &\times \left\{ (x - a_1)^2 \sqrt[4]{185|f'(a_1)|^q + 58|f'(x)|^q} + (a_2 - x)^2 \sqrt[4]{195|f'(x)|^q + 48|f'(a_2)|^q} \right\}. \end{aligned} \quad (2.35)$$

Corollary 2.22. Taking $\lambda = 0$ and $\varphi(t) = t$ in Theorem 2.15, we have

$$\begin{aligned} |T_{f, \Delta_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\eta^{\frac{q+1}{q}}(x, ma_1)\eta(a_2, ma_1)} \\ &\times \left[B_1^g(x; 1) \right]^{1-\frac{1}{q}} \sqrt[4]{\left[B_1^g(x; 1)\eta(x, ma_1) - C_1^g(x) \right] |f'(ma_1)|^q + C_1^g(x) |f'(x)|^q} \\ &\quad + \frac{1}{\eta^{\frac{q+1}{q}}(a_2, mx)\eta(a_2, ma_1)} \left[B_2^g(x; 1) \right]^{1-\frac{1}{q}} \\ &\quad \times \sqrt[4]{\left[B_2^g(x; 1)\eta(a_2, mx) - E_1^g(x) \right] |f'(mx)|^q + E_1^g(x) |f'(a_2)|^q}, \end{aligned} \quad (2.36)$$

where

$$C_1^g(x) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} (t - ma_1)(g(t) - g(ma_1)) dt, \quad (2.37)$$

$$E_1^g(x) = \int_{mx}^{mx + \eta(a_2, mx)} (t - mx)(g(mx + \eta(a_2, mx)) - g(t)) dt, \quad (2.38)$$

and $B_1^g(x; 1)$, $B_2^g(x; 1)$ are defined as in Corollary 2.10 for value $p = 1$.

Corollary 2.23. Taking $\lambda = 0$ and $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$ in Theorem 2.15, we get

$$\begin{aligned} |T_{f, \Lambda_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\eta^{\frac{q+1}{q}}(x, ma_1)\eta(a_2, ma_1)} \\ &\times \left[B_3^g(x; 1, \alpha) \right]^{1-\frac{1}{q}} \\ &\times \sqrt[q]{\left[B_3^g(x; 1, \alpha)\eta(x, ma_1) - C_1^g(x, \alpha) \right] |f'(ma_1)|^q + C_1^g(x, \alpha) |f'(x)|^q} \\ &+ \frac{1}{\eta^{\frac{q+1}{q}}(a_2, mx)\eta(a_2, ma_1)} \left[B_4^g(x; 1, \alpha) \right]^{1-\frac{1}{q}} \\ &\times \sqrt[q]{\left[B_4^g(x; 1, \alpha)\eta(a_2, mx) - E_1^g(x, \alpha) \right] |f'(mx)|^q + E_1^g(x, \alpha) |f'(a_2)|^q}, \end{aligned} \quad (2.39)$$

where

$$C_1^g(x, \alpha) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} (t - ma_1) [g(t) - g(ma_1)]^\alpha dt, \quad (2.40)$$

$$E_1^g(x, \alpha) = \int_{mx}^{mx + \eta(a_2, mx)} (t - mx) [g(mx + \eta(a_2, mx)) - g(t)]^\alpha dt, \quad (2.41)$$

and $B_3^g(x; 1, \alpha)$, $B_4^g(x; 1, \alpha)$ are defined as in Corollary 2.11 for value $p = 1$.

Corollary 2.24. Taking $\lambda = 0$ and $\varphi(t) = \frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ in Theorem 2.15, we obtain

$$\begin{aligned} |T_{f, \Lambda_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\eta^{\frac{q+1}{q}}(x, ma_1)\eta(a_2, ma_1)} \\ &\times \left[B_5^g(x; 1, \alpha, k) \right]^{1-\frac{1}{q}} \\ &\times \sqrt[q]{\left[B_5^g(x; 1, \alpha, k)\eta(x, ma_1) - C_1^g(x, \alpha, k) \right] |f'(ma_1)|^q + C_1^g(x, \alpha, k) |f'(x)|^q} \\ &+ \frac{1}{\eta^{\frac{q+1}{q}}(a_2, mx)\eta(a_2, ma_1)} \left[B_6^g(x; 1, \alpha, k) \right]^{1-\frac{1}{q}} \\ &\times \sqrt[q]{\left[B_6^g(x; 1, \alpha, k)\eta(a_2, mx) - E_1^g(x, \alpha, k) \right] |f'(mx)|^q + E_1^g(x, \alpha, k) |f'(a_2)|^q}, \end{aligned} \quad (2.42)$$

where

$$C_1^g(x, \alpha, k) = \int_{ma_1}^{ma_1 + \eta(x, ma_1)} (t - ma_1) [g(t) - g(ma_1)]^{\frac{\alpha}{k}} dt, \quad (2.43)$$

$$E_1^g(x, \alpha, k) := \int_{mx}^{mx + \eta(a_2, mx)} (t - mx) [g(mx + \eta(a_2, mx)) - g(t)]^{\frac{\alpha}{k}} dt, \quad (2.44)$$

and $B_5^g(x; 1, \alpha, k)$, $B_6^g(x; 1, \alpha, k)$ are defined as in Corollary 2.12 for value $p = 1$.

Corollary 2.25. Taking $\lambda = 0$, $\varphi_g(x, t) = t(g(ma_1 + \eta(x, ma_1)) - t)^{\alpha-1}$ in $[0, t]$ and $\varphi_g(x, t) = t(g(mx + \eta(a_2, mx)) - t)^{\alpha-1}$ in $[t, 1]$ in Theorem 2.15, we have

$$\begin{aligned} |T_{f, \Lambda_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{\eta^{\frac{q+1}{q}}(x, ma_1)\eta(a_2, ma_1)} \\ &\times \left[B_7^g(x; 1, \alpha) \right]^{1-\frac{1}{q}} \sqrt[q]{\left[B_7^g(x; 1, \alpha)\eta(x, ma_1) - C_1^g(x) \right] |f'(ma_1)|^q + C_1^g(x) |f'(x)|^q} \\ &+ \frac{1}{\eta^{\frac{q+1}{q}}(a_2, mx)\eta(a_2, ma_1)} \left[B_8^g(x; 1, \alpha) \right]^{1-\frac{1}{q}} \\ &\times \sqrt[q]{\left[B_8^g(x; 1, \alpha)\eta(a_2, mx) - L_2^g(x, \alpha) \right] |f'(mx)|^q + L_2^g(x, \alpha) |f'(a_2)|^q}, \end{aligned} \quad (2.45)$$

where

$$L_2^g(x, \alpha) = \int_{mx}^{mx+\eta(a_2, mx)} (t - mx) [g^\alpha(mx + \eta(a_2, mx)) - g^\alpha(t)] dt, \quad (2.46)$$

and $B_7^g(x; 1, \alpha)$, $B_8^g(x; 1, \alpha)$ are defined as in Corollary 2.13 for value $p = 1$ and $C_1^g(x)$ is defined as in Corollary 2.22.

Corollary 2.26. Taking $\lambda = 0$ and $\varphi(t) = \frac{t}{\alpha} \exp(-At)$, where $A = \frac{1-\alpha}{\alpha}$ in Theorem 2.15, we get

$$\begin{aligned} |T_{f, \Lambda_m^g, \Delta_m^g}(0; x, a_1, a_2)| &\leq \frac{1}{(1-\alpha)\eta^{\frac{q+1}{q}}(x, ma_1)\eta(a_2, ma_1)} \\ &\times \left\{ \left[B_9^g(x; 1, A) \right]^{1-\frac{1}{q}} \sqrt[q]{L_3^g(x, A) |f'(ma_1)|^q + L_4^g(x, A) |f'(x)|^q} \right. \\ &+ \frac{1}{(1-\alpha)\eta^{\frac{q+1}{q}}(a_2, mx)\eta(a_2, ma_1)} \\ &\left. \times \left[B_{10}^g(x; 1, A) \right]^{1-\frac{1}{q}} \sqrt[q]{L_5^g(x, A) |f'(mx)|^q + L_6^g(x, A) |f'(a_2)|^q} \right\}, \end{aligned} \quad (2.47)$$

where

$$L_3^g(x, A) = \int_{ma_1}^{ma_1+\eta(x, ma_1)} (ma_1 + \eta(x, ma_1) - t) \left\{ 1 - \exp[A(g(ma_1) - g(t))] \right\} dt, \quad (2.48)$$

$$L_4^g(x, A) = \int_{ma_1}^{ma_1+\eta(x, ma_1)} (t - ma_1) \left\{ 1 - \exp[A(g(ma_1) - g(t))] \right\} dt, \quad (2.49)$$

$$\begin{aligned} L_5^g(x, A) &= \int_{mx}^{mx+\eta(a_2, mx)} (mx + \eta(a_2, mx) - t) \\ &\times \left\{ 1 - \exp[A(g(t) - g(mx + \eta(a_2, mx)))] \right\} dt, \end{aligned} \quad (2.50)$$

$$L_6^g(x, A) = \int_{mx}^{mx+\eta(a_2, mx)} (t - mx) \left\{ 1 - \exp[A(g(t) - g(mx + \eta(a_2, mx)))] \right\} dt, \quad (2.51)$$

and $B_9^g(x; 1, A)$, $B_{10}^g(x; 1, A)$ are defined as in Corollary 2.14 for value $p = 1$.

Remark 2.27. Applying our Theorems 2.3 and 2.15 for special values of parameter $\lambda \in [0, 1]$, for appropriate choices of function $g(t) = t$; $g(t) = \ln t$, $\forall t > 0$; $g(t) = e^t$, etc., where $\varphi(t) = t$, $\frac{t^\alpha}{\Gamma(\alpha)}$, $\frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$; $\varphi_g(t) = t(g(a_2) - t)^{\alpha-1}$ for $\alpha \in (0, 1)$; $\varphi(t) = \frac{t}{\alpha} \exp \left[(-\frac{1-\alpha}{\alpha}) t \right]$ for $\alpha \in (0, 1)$, such that $|f'|^q$ to be preinvex (or convex in special case), we can deduce some new general fractional integral inequalities. The details are left to the interested reader.

3. APPLICATIONS

Consider the following special means for different positive real numbers a_1 and a_2 as follows:

(1) The arithmetic mean:

$$A(a_1, a_2) = \frac{a_1 + a_2}{2},$$

(2) The harmonic mean:

$$H(a_1, a_2) = \frac{2}{\frac{1}{a_1} + \frac{1}{a_2}},$$

(3) The logarithmic mean:

$$L(a_1, a_2) = \frac{a_2 - a_1}{\ln a_2 - \ln a_1},$$

(4) The generalized log-mean:

$$L_r(a_1, a_2) = \left[\frac{a_2^{r+1} - a_1^{r+1}}{(r+1)(a_2 - a_1)} \right]^{\frac{1}{r}}; \quad r \in \mathbb{Z} \setminus \{-1, 0\}.$$

Now, using the theory results in section 2, we give some applications to special means for positive different real numbers.

Proposition 3.1. *Let $0 < a_1 < a_2$. Then for $r \in \mathbb{N}$ and $r \geq 2$, where $q > 1$ and $p^{-1} + q^{-1} = 1$, the following inequality holds:*

$$\begin{aligned} & \left| A^r(a_1, a_2) - L_r^r(a_1, a_2) \right| \leq \frac{r(a_2 - a_1)}{4\sqrt[p+1]{p+1}} \\ & \times \left\{ \sqrt[q]{A \left(a_1^{q(r-1)}, \left(\frac{a_1 + a_2}{2} \right)^{q(r-1)} \right)} + \sqrt[q]{A \left(\left(\frac{a_1 + a_2}{2} \right)^{q(r-1)}, a_2^{q(r-1)} \right)} \right\}. \end{aligned} \quad (3.1)$$

Proof. Taking $m = 1$, $\lambda = 0$, $x = \frac{a_1 + a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = t^r$ and $g(t) = \varphi(t) = t$, in Theorem 2.3, one can obtain the result immediately. \square

Proposition 3.2. *Let $0 < a_1 < a_2$. Then for $r \in \mathbb{N}$ and $r \geq 2$, where $q > 1$ and $p^{-1} + q^{-1} = 1$, the following inequality holds:*

$$\begin{aligned} & \left| A(a_1^r, a_2^r) - L_r^r(a_1, a_2) \right| \leq \frac{r(a_2 - a_1)}{4\sqrt[p+1]{p+1}} \\ & \times \left\{ \sqrt[q]{A \left(a_1^{q(r-1)}, \left(\frac{a_1 + a_2}{2} \right)^{q(r-1)} \right)} + \sqrt[q]{A \left(\left(\frac{a_1 + a_2}{2} \right)^{q(r-1)}, a_2^{q(r-1)} \right)} \right\}. \end{aligned} \quad (3.2)$$

Proof. Taking $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = t^r$ and $g(t) = \varphi(t) = t$, in Theorem 2.3, one can obtain the result immediately. \square

Proposition 3.3. *Let $0 < a_1 < a_2$. Then for $q > 1$ and $p^{-1}+q^{-1} = 1$, the following inequality holds:*

$$\left| \frac{1}{A(a_1, a_2)} - \frac{1}{L(a_1, a_2)} \right| \leq \frac{(a_2 - a_1)}{4\sqrt[2]{p+1}} \quad (3.3)$$

$$\times \left\{ \frac{1}{\sqrt[q]{H\left(a_1^{2q}, \left(\frac{a_1+a_2}{2}\right)^{2q}\right)}} + \frac{1}{\sqrt[q]{H\left(\left(\frac{a_1+a_2}{2}\right)^{2q}, a_2^{2q}\right)}} \right\}.$$

Proof. Taking $m = 1$, $\lambda = 0$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = \frac{1}{t}$ and $g(t) = \varphi(t) = t$, in Theorem 2.3, one can obtain the result immediately. \square

Proposition 3.4. *Let $0 < a_1 < a_2$. Then for $q > 1$ and $p^{-1}+q^{-1} = 1$, the following inequality holds:*

$$\left| \frac{1}{H(a_1, a_2)} - \frac{1}{L(a_1, a_2)} \right| \leq \frac{(a_2 - a_1)}{4\sqrt[2]{p+1}} \quad (3.4)$$

$$\times \left\{ \frac{1}{\sqrt[q]{H\left(a_1^{2q}, \left(\frac{a_1+a_2}{2}\right)^{2q}\right)}} + \frac{1}{\sqrt[q]{H\left(\left(\frac{a_1+a_2}{2}\right)^{2q}, a_2^{2q}\right)}} \right\}.$$

Proof. Taking $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = \frac{1}{t}$ and $g(t) = \varphi(t) = t$, in Theorem 2.3, one can obtain the result immediately. \square

Proposition 3.5. *Let $0 < a_1 < a_2$. Then for $r \in \mathbb{N}$ and $r \geq 2$, where $q \geq 1$, the following inequality holds:*

$$\left| A^r(a_1, a_2) - L_r^r(a_1, a_2) \right| \leq \sqrt{\frac{2}{3}} \frac{r(a_2 - a_1)}{8} \quad (3.5)$$

$$\times \left\{ \sqrt[q]{A\left(a_1^{q(r-1)}, 2\left(\frac{a_1+a_2}{2}\right)^{q(r-1)}\right)} + \sqrt[q]{A\left(2\left(\frac{a_1+a_2}{2}\right)^{q(r-1)}, a_2^{q(r-1)}\right)} \right\}.$$

Proof. Taking $m = 1$, $\lambda = 0$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = t^r$ and $g(t) = \varphi(t) = t$, in Theorem 2.15, one can obtain the result immediately. \square

Proposition 3.6. *Let $0 < a_1 < a_2$. Then for $r \in \mathbb{N}$ and $r \geq 2$, where $q \geq 1$, the following inequality holds:*

$$\left| A(a_1^r, a_2^r) - L_r^r(a_1, a_2) \right| \leq \sqrt{\frac{2}{3}} \frac{r(a_2 - a_1)}{8} \quad (3.6)$$

$$\times \left\{ \sqrt[q]{A\left(2a_1^{q(r-1)}, \left(\frac{a_1+a_2}{2}\right)^{q(r-1)}\right)} + \sqrt[q]{A\left(\left(\frac{a_1+a_2}{2}\right)^{q(r-1)}, 2a_2^{q(r-1)}\right)} \right\}.$$

Proof. Taking $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = t^r$ and $g(t) = \varphi(t) = t$, in Theorem 2.15, one can obtain the result immediately. \square

Proposition 3.7. *Let $0 < a_1 < a_2$. Then for $q \geq 1$, the following inequality holds:*

$$\begin{aligned} \left| \frac{1}{A(a_1, a_2)} - \frac{1}{L(a_1, a_2)} \right| &\leq \sqrt[q]{\frac{4}{3}} \frac{(a_2 - a_1)}{8} \\ &\times \left\{ \frac{1}{\sqrt[q]{H\left(2a_1^{2q}, \left(\frac{a_1+a_2}{2}\right)^{2q}\right)}} + \frac{1}{\sqrt[q]{H\left(\left(\frac{a_1+a_2}{2}\right)^{2q}, 2a_2^{2q}\right)}} \right\}. \end{aligned} \quad (3.7)$$

Proof. Taking $m = 1$, $\lambda = 0$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = \frac{1}{t}$ and $g(t) = \varphi(t) = t$, in Theorem 2.15, one can obtain the result immediately. \square

Proposition 3.8. *Let $0 < a_1 < a_2$. Then for $q \geq 1$, the following inequality holds:*

$$\begin{aligned} \left| \frac{1}{H(a_1, a_2)} - \frac{1}{L(a_1, a_2)} \right| &\leq \sqrt[q]{\frac{4}{3}} \frac{(a_2 - a_1)}{8} \\ &\times \left\{ \frac{1}{\sqrt[q]{H\left(a_1^{2q}, 2\left(\frac{a_1+a_2}{2}\right)^{2q}\right)}} + \frac{1}{\sqrt[q]{H\left(2\left(\frac{a_1+a_2}{2}\right)^{2q}, a_2^{2q}\right)}} \right\}. \end{aligned} \quad (3.8)$$

Proof. Taking $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$, $f(t) = \frac{1}{t}$ and $g(t) = \varphi(t) = t$, in Theorem 2.15, one can obtain the result immediately. \square

Remark 3.9. Applying our Theorems 2.3 and 2.15 for special values of parameter $\lambda \in [0, 1]$, for appropriate choices of function $g(t) = t$; $g(t) = \ln t$, $\forall t > 0$; $g(t) = e^t$, etc., where $\varphi(t) = t$, $\frac{t^\alpha}{\Gamma(\alpha)}$, $\frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$; $\varphi_g(t) = t(g(a_2) - t)^{\alpha-1}$ for $\alpha \in (0, 1)$; $\varphi(t) = \frac{t}{\alpha} \exp\left[-\left(\frac{1-\alpha}{\alpha}\right)t\right]$ for $\alpha \in (0, 1)$, such that $|f'|^q$ to be preinvex (or convex in the special case), we can deduce some new general fractional integral inequalities using above special means (and other special means). The details are left to the interested reader.

Next, we provide some new error estimates for the midpoint and trapezium quadrature formula. Let Q be the partition of the points $a_1 = x_0 < x_1 < \dots < x_k = a_2$ of the interval $[a_1, a_2]$. Let consider the following quadrature formula:

$$\int_{a_1}^{a_2} f(x)dx = M(f, Q) + E(f, Q), \quad \int_{a_1}^{a_2} f(x)dx = T(f, Q) + E^*(f, Q),$$

where

$$M(f, Q) = \sum_{i=0}^{k-1} f\left(\frac{x_i + x_{i+1}}{2}\right)(x_{i+1} - x_i)$$

and

$$T(f, Q) = \sum_{i=0}^{k-1} \frac{f(x_i) + f(x_{i+1})}{2} (x_{i+1} - x_i)$$

are the midpoint and trapezium version and $E(f, Q)$, $E^*(f, Q)$ are denote their associated approximation errors.

Proposition 3.10. *Let $f : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function on (a_1, a_2) , where $a_1 < a_2$. If $|f'|^q$ is convex on $[a_1, a_2]$ for $q > 1$ and $p^{-1} + q^{-1} = 1$, then the following inequality holds:*

$$\begin{aligned} |E(f, Q)| &\leq \frac{1}{4\sqrt[q]{2}\sqrt[p]{p+1}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q} + \sqrt[q]{\left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned} \quad (3.9)$$

Proof. Applying Theorem 2.3 for $m = 1$, $\lambda = 0$, $x = \frac{a_1 + a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ on the subintervals $[x_i, x_{i+1}]$ ($i = 0, \dots, k-1$) of the partition Q , we have

$$\begin{aligned} \left| f\left(\frac{x_i + x_{i+1}}{2}\right) - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right| &\leq \frac{(x_{i+1} - x_i)}{4\sqrt[q]{2}\sqrt[p]{p+1}} \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q} + \sqrt[q]{\left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned} \quad (3.10)$$

Hence from (3.10), we get

$$\begin{aligned} |E(f, Q)| &= \left| \int_{a_1}^{a_2} f(x) dx - M(f, Q) \right| \\ &\leq \left| \sum_{i=0}^{k-1} \left\{ \int_{x_i}^{x_{i+1}} f(x) dx - f\left(\frac{x_i + x_{i+1}}{2}\right) (x_{i+1} - x_i) \right\} \right| \\ &\leq \sum_{i=0}^{k-1} \left| \left\{ \int_{x_i}^{x_{i+1}} f(x) dx - f\left(\frac{x_i + x_{i+1}}{2}\right) (x_{i+1} - x_i) \right\} \right| \\ &\leq \frac{1}{4\sqrt[q]{2}\sqrt[p]{p+1}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q} + \sqrt[q]{\left|f'\left(\frac{x_i + x_{i+1}}{2}\right)\right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned}$$

The proof of Proposition 3.10 is completed. \square

Proposition 3.11. *Let $f : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function on (a_1, a_2) , where $a_1 < a_2$. If $|f'|^q$ is convex on $[a_1, a_2]$ for $q \geq 1$, then the following inequality holds:*

$$|E(f, Q)| \leq \frac{1}{8\sqrt[q]{3}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \quad (3.11)$$

$$\times \left\{ \sqrt[q]{|f'(x_i)|^q + 2 \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q} + \sqrt[q]{2 \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + |f'(x_{i+1})|^q} \right\}.$$

Proof. The proof is analogous as to that of Proposition 3.10 taking $m = 1$, $\lambda = 0$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ using Theorem 2.15. \square

Proposition 3.12. *Let $f : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function on (a_1, a_2) , where $a_1 < a_2$. If $|f'|^q$ is convex on $[a_1, a_2]$ for $q > 1$ and $p^{-1} + q^{-1} = 1$, then the following inequality holds:*

$$\begin{aligned} |E^*(f, Q)| &\leq \frac{1}{4\sqrt[4]{2}\sqrt[p+1]{p+1}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q} + \sqrt[q]{\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned} \quad (3.12)$$

Proof. Applying Theorem 2.3 for $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ on the subintervals $[x_i, x_{i+1}]$ ($i = 0, \dots, k-1$) of the partition Q , we have

$$\begin{aligned} \left| \frac{f(x_i) + f(x_{i+1})}{2} - \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx \right| &\leq \frac{(x_{i+1} - x_i)}{4\sqrt[4]{2}\sqrt[p+1]{p+1}} \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q} + \sqrt[q]{\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned} \quad (3.13)$$

Hence from (3.13), we get

$$\begin{aligned} |E^*(f, Q)| &= \left| \int_{a_1}^{a_2} f(x) dx - T(f, Q) \right| \\ &\leq \left| \sum_{i=0}^{k-1} \left\{ \int_{x_i}^{x_{i+1}} f(x) dx - \frac{f(x_i) + f(x_{i+1})}{2} (x_{i+1} - x_i) \right\} \right| \\ &\leq \sum_{i=0}^{k-1} \left| \left\{ \int_{x_i}^{x_{i+1}} f(x) dx - \frac{f(x_i) + f(x_{i+1})}{2} (x_{i+1} - x_i) \right\} \right| \\ &\leq \frac{1}{4\sqrt[4]{2}\sqrt[p+1]{p+1}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \\ &\times \left\{ \sqrt[q]{|f'(x_i)|^q + \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q} + \sqrt[q]{\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + |f'(x_{i+1})|^q} \right\}. \end{aligned}$$

The proof of Proposition 3.12 is completed. \square

Proposition 3.13. *Let $f : [a_1, a_2] \rightarrow \mathbb{R}$ be a differentiable function on (a_1, a_2) , where $a_1 < a_2$. If $|f'|^q$ is convex on $[a_1, a_2]$ for $q \geq 1$, then the following inequality holds:*

$$|E^*(f, Q)| \leq \frac{1}{8\sqrt[4]{3}} \times \sum_{i=0}^{k-1} (x_{i+1} - x_i)^2 \quad (3.14)$$

$$\times \left\{ \sqrt[q]{2|f'(x_i)|^q + \left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q} + \sqrt[q]{\left| f' \left(\frac{x_i + x_{i+1}}{2} \right) \right|^q + 2|f'(x_{i+1})|^q} \right\}.$$

Proof. The proof is analogous as to that of Proposition 3.12 taking $m = 1$, $\lambda = 1$, $x = \frac{a_1+a_2}{2}$, $\eta(x, ma_1) = x - ma_1$, $\eta(a_2, mx) = a_2 - mx$, $\eta(a_2, ma_1) = a_2 - ma_1$ and $g(t) = \varphi(t) = t$ using Theorem 2.15. \square

Remark 3.14. Applying our Theorems 2.3 and 2.15, where $m = 1$, for special values of parameter $\lambda \in [0, 1]$, for appropriate choices of function $g(t) = t$; $g(t) = \ln t$, $\forall t > 0$; $g(t) = e^t$, etc., where $\varphi(t) = t$, $\frac{t^\alpha}{\Gamma(\alpha)}$, $\frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$; $\varphi_g(t) = t(g(a_2) - t)^{\alpha-1}$ for $\alpha \in (0, 1)$; $\varphi(t) = \frac{t}{\alpha} \exp \left[\left(-\frac{1-\alpha}{\alpha} \right) t \right]$ for $\alpha \in (0, 1)$, such that $|f'|^q$ to be convex, we can deduce some new bounds for the midpoint and trapezium quadrature formula using above ideas and techniques. The details are left to the interested reader.

Acknowledgements

The authors would like to thank the honorable referee for valuable comments and suggestions in order to improve our manuscript.

REFERENCES

- [1] R. P. Agarwal, M. J. Luo and R. K. Raina, On Ostrowski type inequalities, *Fasc. Math.*, **204** (2016), 5–27.
- [2] M. Ahmadmir and R. Ullah, Some inequalities of Ostrowski and Grüss type for triple integrals on time scales, *Tamkang J. Math.*, **42**(4) (2011), 415–426.
- [3] M. Alomari, M. Darus, S. S. Dragomir and P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, *Appl. Math. Lett.*, **23** (2010), 1071–1076.
- [4] S. M. Aslani, M. R. Delavar and S. M. Vaezpour, Inequalities of Fejér type related to generalized convex functions with applications, *Int. J. Anal. Appl.*, **16**(1) (2018), 38–49.
- [5] F. X. Chen and S. H. Wu, Several complementary inequalities to inequalities of Hermite–Hadamard type for s -convex functions, *J. Nonlinear Sci. Appl.*, **9**(2) (2016), 705–716.
- [6] Y. M., Chu, M. A. Khan, T. U. Khan and T. Ali, Generalizations of Hermite–Hadamard type inequalities for MT -convex functions, *J. Nonlinear Sci. Appl.*, **9**(5) (2016), 4305–4316.
- [7] M. R. Delavar and S. S. Dragomir, On η -convexity, *Math. Inequal. Appl.*, **20** (2017), 203–216.
- [8] M. R. Delavar and M. De La Sen, Some generalizations of Hermite–Hadamard type inequalities, *SpringerPlus*, **5**(1661) (2016).
- [9] S. S. Dragomir, On the Ostrowski’s integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, **1**(2) (1998).
- [10] S. S. Dragomir, The Ostrowski integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, **38** (1999), 33–37.
- [11] S. S. Dragomir, Ostrowski-type inequalities for Lebesgue integral: A survey of recent results, *Aust. J. Math. Anal. Appl.*, **14**(1) (2017), 1–287.
- [12] S. S. Dragomir, Further Ostrowski and trapezoid type inequalities for the generalized Riemann–Liouville fractional integrals of functions with bounded variation, *RGMIA Res. Rep. Coll.*, **20** (2017), Art. 84, <http://rgmia.org/papers/v20/v20a84.pdf>.
- [13] S. S. Dragomir, Ostrowski and trapezoid type inequalities for the generalized k - g -fractional integrals of functions with bounded variation, *RGMIA Res. Rep. Coll.*, **20** (2017), Art. 111, <http://rgmia.org/papers/v20/v20a111.pdf>.
- [14] S. S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and trapezoidal formula, *Appl. Math. Lett.*, **11**(5) (1998), 91–95.

- [15] S. S. Dragomir and S. Wang, An inequality of Ostrowski–Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules, *Comput. Math. Appl.*, **13**(11) (1997), 15–20.
- [16] S. S. Dragomir and S. Wang, A new inequality of Ostrowski’s type in L_1 –norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. Math.*, **28** (1997), 239–244.
- [17] T. S. Du, J. G. Liao and Y. J. Li, Properties and integral inequalities of Hadamard–Simpson type for the generalized (s, m) –preinvex functions, *J. Nonlinear Sci. Appl.*, **9** (2016), 3112–3126.
- [18] G. Farid, Existence of an integral operator and its consequences in fractional and conformable integrals, *Open J. Math. Sci.*, **3** (2019), 210–216.
- [19] G. Farid, Some new Ostrowski type inequalities via fractional integrals, *Int. J. Anal. App.*, **14**(1) (2017), 64–68.
- [20] J. Hristov, Response functions in linear viscoelastic constitutive equations and related fractional operators, *Math. Model. Nat. Phenom.*, **14**(3) (2019), 1–34.
- [21] M. Jleli and B. Samet, On Hermite–Hadamard type inequalities via fractional integral of a function with respect to another function, *J. Nonlinear Sci. Appl.*, **9** (2016), 1252–1260.
- [22] A. Kashuri and R. Liko, Some new Hermite–Hadamard type inequalities and their applications, *Stud. Sci. Math. Hung.*, **56**(1) (2019), 103–142.
- [23] A. Kashuri and R. Liko, Ostrowski type fractional integral inequalities for generalized (s, m, φ) –preinvex functions, *Aust. J. Math. Anal. Appl.*, **13**(1) (2016), Art. 16, 1–11.
- [24] A. Kashuri and R. Liko, Generalizations of Hermite–Hadamard and Ostrowski type inequalities for MT_m –preinvex functions, *Proyecciones*, **36**(1) (2017), 45–80.
- [25] U. N. Katugampola, New approach to a generalized fractional integral, *Appl. Math. Comput.*, **218**(3) (2011), 860–865.
- [26] M. A. Khan, Y. M. Chu, A. Kashuri and R. Liko, Hermite–Hadamard type fractional integral inequalities for $MT_{(r;g,m,\varphi)}$ –preinvex functions, *J. Comput. Anal. Appl.*, **26**(8) (2019), 1487–1503.
- [27] M. A. Khan, Y. M. Chu, A. Kashuri, R. Liko and G. Ali, Conformable fractional integrals versions of Hermite–Hadamard inequalities and their generalizations, *J. Funct. Spaces*, (2018), Art. ID 6928130, pp. 9.
- [28] A. A. Kilbas, O. I. Marichev and S. G. Samko, Fractional Integrals and Derivatives. Theory and Applications, *Gordon and Breach, Switzerland*, (1993).
- [29] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, *Elsevier Science B.V., Amsterdam*, (2006).
- [30] R. Khalil, M. A. Horani, A. Yousef and M. Sababheh, A new definition of fractional derivatives, *J. Comput. Appl. Math.*, **264** (2014), 65–70.
- [31] M. Kirane and B. T. Torebek, Hermite–Hadamard, Hermite–Hadamard–Fejér, Dragomir–Agarwal and Pachpatte type inequalities for convex functions via fractional integrals, *arXiv:1701.00092*.
- [32] W. J. Liu, Some Simpson type inequalities for h –convex and (α, m) –convex functions, *J. Comput. Anal. Appl.*, **16**(5) (2014), 1005–1012.
- [33] W. Liu, W. Wen and J. Park, Hermite–Hadamard type inequalities for MT –convex functions via classical integrals and fractional integrals, *J. Nonlinear Sci. Appl.*, **9** (2016), 766–777.
- [34] W. Liu, W. Wen and J. Park, Ostrowski type fractional integral inequalities for MT –convex functions, *Miskolc Math. Notes*, **16**(1) (2015), 249–256.
- [35] Z. Liu, Some Ostrowski–Grüss type inequalities and applications, *Comput. Math. Appl.*, **53** (2007), 73–79.
- [36] Z. Liu, Some companions of an Ostrowski type inequality and applications, *J. Inequal. in Pure and Appl. Math.*, **10**(2) (2009), Art. 52, pp. 12.
- [37] C. Luo, T. S. Du, M. A. Khan, A. Kashuri and Y. Shen, Some k –fractional integrals inequalities through generalized $\lambda_{\phi m}$ – MT –preinvexity, *J. Comput. Anal. Appl.*, **27**(4) (2019), 690–705.
- [38] M. Matloka, Ostrowski type inequalities for functions whose derivatives are h –convex via fractional integrals, *Journal of Scientific Research and Reports*, **3**(12) (2014), 1633–1641.
- [39] M. V. Mihai, Some Hermite–Hadamard type inequalities via Riemann–Liouville fractional calculus, *Tamkang J. Math.*, **44**(4) (2013), 411–416.

- [40] S. Mubeen and G. M. Habibullah, k -Fractional integrals and applications, *Int. J. Contemp. Math. Sci.*, **7** (2012), 89–94.
- [41] O. Omotoyinbo and A. Mogbodemu, Some new Hermite–Hadamard integral inequalities for convex functions, *Int. J. Sci. Innovation Tech.*, **1**(1) (2014), 1–12.
- [42] M. E. Özdemir, S. S. Dragomir and C. Yıldız, The Hadamard's inequality for convex function via fractional integrals, *Acta Math. Sci., Ser. B, Engl. Ed.*, **33**(5) (2013), 153–164.
- [43] M. E. Özdemir, H. Kavurmacı and E. Set, Ostrowski's type inequalities for (α, m) -convex functions, *Kyungpook Math. J.*, **50** (2010), 371–378.
- [44] B. G. Pachpatte, On an inequality of Ostrowski type in three independent variables, *J. Math. Anal. Appl.*, **249** (2000), 583–591.
- [45] B. G. Pachpatte, On a new Ostrowski type inequality in two independent variables, *Tamkang J. Math.*, **32**(1) (2001), 45–49.
- [46] F. Qi and B. Y. Xi, Some integral inequalities of Simpson type for $GA - \epsilon$ -convex functions, *Georgian Math. J.*, **20**(5) (2013), 775–788.
- [47] A. Rafiq, N. A. Mir and F. Ahmad, Weighted Čebyšev–Ostrowski type inequalities, *Applied Math. Mechanics (English Edition)*, **28**(7) (2007), 901–906.
- [48] M. Z. Sarikaya, On the Ostrowski type integral inequality, *Acta Math. Univ. Comenianae*, **79**(1) (2010), 129–134.
- [49] M. Z. Sarikaya and F. Ertuğral, On the generalized Hermite–Hadamard inequalities, <https://www.researchgate.net/publication/321760443>.
- [50] M. Z. Sarikaya and H. Yıldırım, On generalization of the Riesz potential, *Indian Jour. of Math. and Mathematical Sci.*, **3**(2), (2007), 231–235.
- [51] E. Set, M. A. Noor, M. U. Awan and A. Gözpınar, Generalized Hermite–Hadamard type inequalities involving fractional integral operators, *J. Inequal. Appl.*, **169** (2017), 1–10.
- [52] M. Tunç, Ostrowski type inequalities for functions whose derivatives are MT -convex, *J. Comput. Anal. Appl.*, **17**(4) (2014), 691–696.
- [53] N. Ujević, Sharp inequalities of Simpson type and Ostrowski type, *Comput. Math. Appl.*, **48** (2004), 145–151.
- [54] H. Wang, T. S. Du and Y. Zhang, k -fractional integral trapezium-like inequalities through (h, m) -convex and (α, m) -convex mappings, *J. Inequal. Appl.*, **2017**(311) (2017), pp. 20.
- [55] B. Y. Xi and F. Qi, Some integral inequalities of Hermite–Hadamard type for convex functions with applications to means, *J. Funct. Spaces Appl.*, **2012** (2012), Art. ID 980438, pp. 14.
- [56] Ç. Yıldız, M. E. Özdemir and M. Z. Sarikaya, New generalizations of Ostrowski–like type inequalities for fractional integrals, *Kyungpook Math. J.*, **56** (2016), 161–172.
- [57] X. M. Zhang, Y. M. Chu and X. H. Zhang, The Hermite–Hadamard type inequality of GA -convex functions and its applications, *J. Inequal. Appl.*, (2010), Art. ID 507560, pp. 11.
- [58] Y. Zhang, T. S. Du, H. Wang, Y. J. Shen and A. Kashuri, Extensions of different type parameterized inequalities for generalized (m, h) -preinvex mappings via k -fractional integrals, *J. Inequal. Appl.*, **2018**(49) (2018), pp. 30.
- [59] L. Zhongxue, On sharp inequalities of Simpson type and Ostrowski type in two independent variables, *Comput. Math. Appl.*, **56** (2008), 2043–2047.

ARTION KASHURI

DEPARTMENT OF MATHEMATICS, FACULTY OF TECHNICAL SCIENCE, UNIVERSITY “ISMAIL QEMALI”,
VLORA, ALBANIA

E-mail address: artionkashuri@gmail.com

MUHAMMAD AMER LATIF

DEPARTMENT OF BASIC SCIENCES, DEANSHIP OF PREPARATORY YEAR, UNIVERSITY OF HAIL, HAIL
2440, SAUDI ARABIA

E-mail address: m_amer_latif@hotmail.com