

## SYSTEMATICALLY SEVERAL GENERAL CLASSES OF BILATERAL GENERATING FUNCTIONS

LAKSHMI NARAYAN MISHRA, RAKESH KUMAR SINGH, SHIKHA PANDEY

**ABSTRACT.** The object of this paper is to present rather systematically several classes of bilateral generating functions which are applicable to such familiar orthogonal polynomials as the Jacobi, Laguerre, Hermite and Bessel polynomials and indeed also to their numerous interesting unifications and generalizations introduced and studied in the literature. Several remarks, comments and observations, relevant to the present discussion are also included.

### 1. INTRODUCTION

Let  $f_k : \mathbb{C} \rightarrow \mathbb{C}$  be a complex-valued polynomial function of a complex variable  $z (k = 1, \dots, n)$ . Also let

$$M_f = \max_{|z|=1} |f(z)|. \quad (1.1)$$

In terms of degree  $d_k$  of the polynomial  $f_k(z) (k = 1, \dots, n)$ , Rassias [6] gave an interesting two-sided inequality for  $M_{f_1 \dots f_n}$ . Indeed, just as the upper and lower bounds derived earlier by Srivastava and Brenner [7] for a general system of orthogonal polynomials, Rassias's result would apply easily to various members of the family of classical orthogonal polynomial (see, e.g., [1], [5], [13] and [14]) led by the Jacobi polynomials:

$$\begin{aligned} p_n^{(\alpha, \beta)}(z) &= \sum_{k=0}^n \begin{bmatrix} \alpha + n \\ n - k \end{bmatrix} \begin{bmatrix} \beta + n \\ k \end{bmatrix} \left[ \frac{z-1}{2} \right]^k \left[ \frac{z+1}{2} \right]^{n-k} \\ &= \begin{bmatrix} \alpha + n \\ n \end{bmatrix} {}_2F_1 \left[ \begin{matrix} -n, \alpha + \beta + n + 1 \\ \alpha + 1 \end{matrix}; \frac{1-z}{2} \right], \end{aligned} \quad (1.2)$$

where  ${}_2F_1$  denotes the Gaussian hypergeometric function defined, in the special case  $p = 2$  and  $q = 1$ , by the power series:

$${}_pF_q \left[ \begin{matrix} \alpha_1, \dots, \alpha_p \\ \beta_1, \dots, \beta_q \end{matrix}; z \right] = \sum_{m=0}^{\infty} \frac{(\alpha_1)_m, \dots, (\alpha_p)_m}{(\beta_1)_m, \dots, (\beta_q)_m} \frac{z^m}{m!} \quad (1.3)$$

$$(p \leq q, |Z| < \infty; p = q + 1, |Z| < 1; \beta_j \neq 0, -1, -2, \dots (j = 1, \dots, q))$$

2010 *Mathematics Subject Classification.* 33D15, 33D10, 33E05.

*Key words and phrases.* polynomial, bilateral generating functions, non-vanishing function, non-negative function.

Submitted April 12, 2020.

with  $(\lambda)_m = \frac{\Gamma(\lambda+m)}{\Gamma(\lambda)}$ .

Some other members of the family of classical orthogonal polynomials include such special cases of the Jacobi polynomials as the Gegenbauer (or ultraspherical) polynomials, the Legendre (or spherical) polynomials, and the Tchebycheff polynomials of the first and second kinds. Indeed we also have, in the same family the classical Laguerre polynomials:

$$L_n^\alpha(z) = \sum_{k=0}^n \begin{bmatrix} \alpha + n \\ n - k \end{bmatrix} \frac{(-z)^k}{k!} = \begin{bmatrix} \alpha - n \\ n \end{bmatrix} {}_2F_1 \left[ \begin{matrix} -n \\ \alpha + 1 \end{matrix}; z \right], \tag{1.4}$$

the classical Hermite polynomials:

$$\begin{aligned} H_n(z) &= \sum_{k=0}^{[n/2]} (-1)^k \begin{bmatrix} n \\ 2k \end{bmatrix} \frac{(2k)!}{k!} (2z)^{n-2k} \\ &= (2z)^n {}_2F_0 \left[ \begin{matrix} -\frac{1}{2}n, -\frac{1}{2}n + \frac{1}{2} \\ - \end{matrix}; -\frac{1}{2} \right], \end{aligned} \tag{1.5}$$

and the Bessel polynomials (cf. [2] and [3]):

$$y_n(z, \alpha, \beta) = \begin{bmatrix} n \\ k \end{bmatrix} k! \left[ \frac{z}{\beta} \right]^k {}_2F_0 \left[ \begin{matrix} -n, \alpha + n - 1 \\ - \end{matrix}; -\frac{z}{\beta} \right], \tag{1.6}$$

which are all limiting cases of the Jacobi polynomials (see, for details, [13, p. 103, Equation (5,3,4)] and [10, p.99, Problems 35 and 36]).

The importance of a systematic study of various properties and characteristics of the classical Jacobi polynomials (including, example linear, bilinear and bilateral generating functions (cf. [4] and [10]) associated with them) led Srivastava and Popov [12] recently to a class of mixed multilateral generating functions for the Jacobi polynomials, which can indeed be suitable applied to derive numerous further results involving Jacobi polynomials and some of their aforementioned relatives.

A familiar unification of each of the classes of orthogonal polynomial named above, and of their various known generalization studied in the literature (cf., e.g., [10]), is provided by the sequence [11, p.307, Equation(5)]:

$$S_n^{(\alpha, \beta)} [x, a, b, c, d; \gamma, \delta; \omega(x)] = \frac{(ax + b)^{-\alpha} (cx + d)^{-\beta}}{n! \omega(x)} D_x^n \{ (ax + b)^{\gamma+n+b} (cx + d)^{\delta+n+\beta} \omega(x) \} \tag{1.7}$$

$$\left[ D_x = \frac{d}{dx}; \quad n = 0, 1, 2, \dots \right],$$

where the parameters  $a, b, c, d, \alpha, \beta, \gamma, \delta$  are arbitrary constants, real or complex and  $\omega(x)$  is independent of  $n$  and differentiable any number of times. For example, it is easily observed from (1.7) and the known Rodrigues formulas for the Jacobi and Bessel polynomials that

$$S_n^{(\alpha, \beta)} [x, a, -a, c, c; 1, 1; C] = (2ca)^n p_n^{\alpha, \beta}(x) \tag{1.8}$$

for any nonzero constant  $C$ , and

$$S_n^{(\alpha-2, 0)} [x, a, 0, c, D; 2, 0; \exp(-\beta/x)] = \frac{\alpha^{2n} \beta^n}{n!} y_n(x, \alpha, \beta). \tag{1.9}$$

Indeed, in each of these and other cases of reducibility, we can have numerous alternative sets of choices for the various parameters involved.

For the sequence of function defined by (1.7), various classes of linear, bilinear, bilateral, and mixed multilateral generating functions are given in the literature. The most general results of this type for the sequence defined by (1.7) were announced recently by Srivastava and Panda [9]. The object of this paper is to present a detailed demonstration of each of the earlier results and to recall how these general results can indeed be applied to derive the corresponding generating functions for various systems of orthogonal polynomials and for numerous other polynomials associated with them [15]-[28].

## 2. GENERAL BILATERAL GENERATING FUNCTIONS

The main results on bilateral generating functions for the sequences defined by (1.7) are given by Theorem 1, 2 and 3 below (cf.[9]).

**Theorem 1.** *Corresponding to a non-vanishing function  $\Omega_\mu(y_1, \dots, y_s)$  of  $s$  variables  $\Omega_\mu(y_1, \dots, y_s)$  ( $s \geq 1$ ) and of (complex) order  $\mu$ , let*

$$E_{m,p,q}^{(p,q)}[x_1; y_1, \dots, y_s; t] = \sum_{n=0}^{\infty} S_{m+qn}^{(\alpha-pqn, \beta-\sigmaqn)}[x, a, b, c, d; \gamma, \delta; \omega(x)] \cdot \Omega_{\mu+pn}(y_1, \dots, y_s) t^n \quad (a_n \neq 0). \quad (2.1)$$

where  $p$  and  $q$  are positive integers,  $\rho$  and  $\sigma$  are complex parameters. Also, for an arbitrary integers  $m \geq 1$ , let

$$\theta_{n,m,p,q}^{\alpha,\beta,\rho,\sigma}[x; y_1, \dots, y_s; z] = \sum_{k=0}^{[n/q]} \begin{bmatrix} m+n \\ n-qn \end{bmatrix} a_k \cdot S_{m+n}^{(\alpha-\gamma n+\rho qk, \beta-\delta n+\sigma qk)}[x, a, b, c, d; \gamma, \delta; \omega(x)] \cdot \Omega_{\mu+pk}(y_1, \dots, y_s) z^k. \quad (2.2)$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} \theta_{n,m,p,q}^{\alpha,\beta,\rho,\sigma}[x; y_1, \dots, y_s; z] &= [1 + at(ax + b^{y-1})(cx + d)^\delta]^\alpha \\ &\cdot [1 + ct(ax + b)^\gamma (cx + d)^{\delta-1}]^\beta \frac{\omega(\xi)}{\omega(x)} F_{m,p,q}^{(\gamma-\rho, \gamma-\sigma)} \\ &[\xi; y_1, \dots, y_s; zt^q \{1 + at(ax + b)^{\gamma-1}(cx + d)^\delta\}^{(\rho-\gamma)q} \\ &\{1 + ct(ax + b)^\gamma (cx + d)^{\delta-1}\}^{(\sigma-\delta)q}], \end{aligned} \quad (2.3)$$

Provided that each side exist; here, for convenience,

$$\xi \equiv \xi(x, t) = x + t(ax + b)^\gamma (cx + d)^\delta. \quad (2.4)$$

**Theorem 2.** *Under the hypotheses of Theorem 1, let*

$$G_{m,p,q}^{(\rho)}[x; y_1, \dots, y_s; t] = \sum_{n=0}^{\infty} a_n S_{m+qn}^{(\alpha-\rhoqn, \beta)}[x, a, b, c, d; \gamma, \delta, \omega(x)] \cdot \Omega_{\mu+qn}(y_1, \dots, y_s) t^n \quad (a_n \neq 0), \quad (2.5)$$

and define

$$\begin{aligned} \Phi_{n,m,p,q}^{\alpha,\beta,\rho}[x; y_1, \dots, y_s; z] = & \sum_{n=0}^{[n/q]} \begin{bmatrix} m+n \\ n-qk \end{bmatrix} a_k \\ & \cdot S_{m+n}^{(\alpha-\gamma n+\rho qk,\beta)}[x, a, b, c, d; \gamma, 1; \omega(x)] \Omega_{\mu+pk}(y_1, \dots, y_s) z^k. \end{aligned} \tag{2.6}$$

for an arbitrary integer  $m \geq 0$ .

Then

$$\begin{aligned} \sum_{n=0}^{\infty} \Phi_{n,m,p,q}^{\alpha,\beta,\rho}[x; y_1, \dots, y_s; z] t^n = & [1 + t(ad - bc)(ax + b)^{\gamma-1}]^\alpha \\ & \cdot [1 + ct(ax + b)^\gamma]^{-\alpha-\beta-m-1} \frac{\omega(\eta)}{\omega(x)} G_{m,p,q}^{(\gamma-\rho)}[\eta; y_1, \dots, y_s; zt^q \\ & \{t(ad - bc)(ax + b)^{\gamma-1}\}^{(\rho-\sigma)q} \{1 - ct(ax + b)^\gamma\}^{(\gamma-\rho-1)q}], \end{aligned} \tag{2.7}$$

where

$$\eta \equiv \eta(x, t) = [x + dt(ax + b)^\gamma] / [1 - ct(ax + b)^\gamma]. \tag{2.8}$$

**Theorem 3.** Under the hypotheses of Theorem 1, let

$$H_{n,m,p,q}^{(\sigma)}[x; y_1, \dots, y_s; t] = \sum_{n=0}^{\infty} a_n S_{m+qn}^{(\alpha,\beta-\sigma\rho n)}[x, a, b, c, d; \gamma, \delta; \omega(x)] \cdot \Omega_{\mu+pn}(y_1, \dots, y_s) t^n \quad (a_n \neq 0), \tag{2.9}$$

and define

$$\begin{aligned} \psi_{n,m,p,q}^{\alpha,\beta,\rho}[x; y_1, \dots, y_s; z] = & \sum_{k=0}^{[n/q]} \begin{bmatrix} m+n \\ n-qk \end{bmatrix} a_k \\ & \cdot S_{m+n}^{(\alpha,\beta-\delta n+\sigma qk)}[x, a, b, c, d; \delta, 1; \omega(x)] \Omega_{\mu+pk}(y_1, \dots, y_s) z^k, \end{aligned} \tag{2.10}$$

for an arbitrary integer  $m \geq 0$ .

Then

$$\begin{aligned} \sum_{n=0}^{\infty} \Psi_{n,m,p,q}^{\alpha,\beta,\rho}[x; y_1, \dots, y_s; z] t^n = & [1 + t(bc - ad)(cx + d)^{\delta-1}]^\beta \\ & \cdot [1 + at(cx + d)^\delta]^{-\alpha-\beta-m-1} \frac{\omega(\zeta)}{\omega(x)} G_{m,p,q}^{(\delta-\rho)}[\zeta; y_1, \dots, y_s; zt^q \\ & \{1 + t(bc - ad)(cx + d)^{\delta-1}\}^{(\sigma-\gamma)q} \{1 - at(cx + d)^\delta\}], \end{aligned} \tag{2.11}$$

where:

$$\zeta \equiv \zeta(x, t) = [x + bt(cx + b)^\delta] / [1 - at(cx + b)^\delta]. \tag{2.12}$$

**Demonstrations**

The proofs of Theorems 1, 2, and 3 are based rather heavily upon certain generating relationships which we recall here as the following

**Lemma 1.** (Srivastava and Panda [11]; see also Srivastava [7]). Let  $\xi, \eta$  and  $\zeta$  and be defined by Equations (2.4), (2.8) and (2.12), respectively.

Then, for every non-negative integer  $m$ ,

$$\begin{aligned} & \sum_{n=0}^{\infty} \begin{bmatrix} m+n \\ n \end{bmatrix} S_{m+n}^{(\alpha-\gamma n, \beta-\delta n)}[x, a, b, c, d; \gamma, \delta; \omega(x)] t^n \\ &= [1 + at(ax+b)^{\gamma-1}(cx+d)^{\delta-1}]^{\beta} \frac{\omega(\xi)}{\omega(x)} G_m^{(\alpha, \beta)}[\xi; a, b, c, d; \gamma, \delta; \omega(\xi)] \end{aligned} \quad (2.13)$$

$$\begin{aligned} & \sum_{n=0}^{\infty} \begin{bmatrix} m+n \\ n \end{bmatrix} S_{m+n}^{(\alpha-\gamma n, \beta)}[x, a, b, c, d; \gamma, 1; \omega(x)] t^n \\ &= \{1 + t(ad-bc)(ax+b)^{\gamma-1}\}^{\alpha} \\ & \cdot [1 - ct(ax+b)^{\gamma}]^{\alpha-\beta-m-1} \frac{\omega(\eta)}{\omega(x)} S_m^{(\alpha, \beta)}[\eta; a, b, c, d; \gamma, 1; \omega(\eta)], \end{aligned} \quad (2.14)$$

and

$$\begin{aligned} & \sum_{n=0}^{\infty} \begin{bmatrix} m+n \\ n \end{bmatrix} S_{m+n}^{(\alpha-\gamma n, \beta)}[x, a, b, c, d; 1, \delta; \omega(x)] t^n \\ &= \{1 + t(bc-ad)(cx+d)^{\delta-1}\}^{\beta} \\ & \cdot [1 - at(cx+d)^{\delta}]^{\alpha-\beta-m-1} \frac{\omega(\zeta)}{\omega(x)} S_m^{(\alpha, \beta)}[\zeta; a, b, c, d; 1, \delta; \omega(\zeta)]. \end{aligned} \quad (2.15)$$

**Proof of Theorem 1.** Denoting the left side of the assertion (2.3) by  $\Delta$ , and substituting for the  $\theta$ -polynomials from (2.2), we observe that

$$\begin{aligned} \Delta &= \sum_{n=0}^{\infty} t^n \sum_{n=0}^{\lfloor n/q \rfloor} \begin{bmatrix} m+n \\ n-qn \end{bmatrix} a_k S_{m+n}^{(\alpha-\gamma n+pqk, \beta-\delta n+\sigma qk)}[x, a, b, c, d; \gamma, \delta; \omega(x)] \\ & \cdot \Omega_{\mu+pk}(y_1, \dots, y_s) z^k \\ &= \sum_{n=0}^{\infty} \Omega_{\mu+pk}(y_1, \dots, y_s) (zt^q)^k \\ & \cdot \sum_{n=0}^{\infty} \begin{bmatrix} m+qn+n \\ n \end{bmatrix} S_{m+n}^{(\alpha+(\rho-\gamma), qk-\gamma n, \beta+(\sigma-\delta)qk-\delta n)}[x, a, b, c, d; \gamma, \delta; \omega(x)] t^n, \end{aligned} \quad (2.16)$$

where we have inverted the order of the double summation involved. If we now sum the inner series in (2.16) by appealing to the generating relation (2.13), with  $m, \alpha$  and  $\beta$  replaced by  $m+qn, \alpha+(\rho-\gamma)qk$ , and  $\beta+(\rho-\delta)qk$ , respectively

( $k = 0, 1, 2, \dots$ ), we shall find that

$$\begin{aligned} \Delta &= [1 + at(ax + b)^{\gamma-1}(cx + d)^{\delta}]^{\alpha} [1 + ct(ax + b)^{\gamma}(cx + d)^{\delta-1}]^{\beta} \frac{\omega(\xi)}{\omega(x)} \\ &\quad \sum_{n=0}^{\infty} a_n S_{m+n}^{(\alpha+(\rho-\gamma)qk, \beta+(\sigma-\delta)qk)}[\xi; a, b, c, d; \gamma, \delta; \omega(\xi)] \Omega_{\mu+pk}(y_1, \dots, y_s) \\ &\quad \cdot \left[ zt^q [1 + at(ax + b)^{\gamma-1}(cx + d)^{\delta}]^{(\rho-\sigma)q} [1 + ct(ax + b)^{\gamma}(cx + d)^{\delta-1}]^{(\sigma-\delta)q} \right]^k, \end{aligned} \tag{2.17}$$

where  $\xi$  is given by (2.4). Interpretation of the infinite series occurring in (2.18) by means of the definition (2.1) yields the right side of the assertion (2.3).

This evidently completes the proof of Theorem 1 under the assumption that the double series involved in the first two steps of our proof are absolutely convergent. Thus, in general, Theorem 1 holds true for these values of the various parameters and variables involved for which each side of the assertion (2.3) exists.

**Proofs of Theorems 2 and 3.** The proofs of Theorems 2 and 3 run parallel to that of Theorem 1, which is already detailed above. Indeed, instead of (2.13), our proofs of Theorems 2 and 3 would make use of the generating relations (2.14) and (2.15), respectively.

Each of the cases of reducibility of the sequence defined by (1.7), which are illustrated by (1.8) and (1.9) and also described fairly adequately by Srivastava and Panda [11, p. 308], can indeed be shown to lead from Theorems 1, 2, and 3 to the corresponding generating functions for simpler (and, quite frequently, more familiar) sequences of special functions and polynomials (see also Remark 3 below). For instance, in one of such circumstances, Theorem 1 will readily yield the main result of Srivastava and Popov [12].

We conclude this section by recalling the following relevant remarks concerning Theorems 1, 2, and 3 (cf. [9]).

**Remark 1.** Since [11, p. 308, Equation (10)]

$$S_m^{\alpha, \beta}[x, a, b, c, d; 1, \delta; \omega(x)] = [ab^{\gamma+\delta-1}c^{\gamma-1}d^{\delta-1}]^{-n} S_m^{\alpha, \beta} \left[ \frac{bcx}{ad}, a, b, a^2d, b^2c; \delta, \gamma; \omega \left( \frac{bcx}{ad} \right) \right]. \tag{2.18}$$

The generating relationships (2.14) and (2.15), and hence also Theorem 2 and Theorem 3, are essentially equivalent.

**Remark 2.** For  $p = \sigma = 0$ , the assertions (2.3), (2.7), and (2.11) would immediately yield certain bilateral generating functions due to Srivastava ([7, Part II, pp. 243-244, Corollaries 19, 20, and 21]; see also [10, pp. 435-436]).

**Remark 3.** By appealing to the known relationships between the sequence (1.7) and its numerous special cases (cf. [11, p. 308]), it is not difficult to derive, from Theorems 1, 2, and 3, analogous results on generating functions for such familiar orthogonal polynomials as Jacobi, Laguerre, Hermite, and Bessel polynomials, and

also for the various interesting generalizations of these polynomials studied in the literature (cf. [3], [10], and [13]). The details involved may be omitted.

**Conflict of Interest** The authors declare that there is no conflict of interests regarding publication of this manuscript.

#### REFERENCES

- [1] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, London, and Paris, 1978.
- [2] E. Grosswald, *Bessel Polynomials*, Springer-Verlag, Berlin, Heidelberg, and New York, 1978.
- [3] H. I. Krall, and O. Frink, A new class of orthogonal polynomials: The Bessel polynomials, *Trans. Amer. lath. Soc.* 65,100-115, 1949.
- [4] E. B. McBride, *Obtaining Generating Functions* (Springer Tracts in Natural Philosophy 21), Springer-Verlag, New York, Heidelberg, and Berlin, 1971.
- [5] E. D. Rainville, *Special Functions*, Macmillan, New York, Reprinted by Chelsea Publishing Company, Bronx, New York, 1971.
- [6] Th. M. Rassias, A new inequality for complex—valued polynomial functions, *Proc. Amer. lath. Soc.* 97,296-298, 1986.
- [7] H. M. Srivastava, Some bilateral generating functions for a certain class of special functions. I and n, *Nederl. Akad. Yetensch. Indag. Math.* 42, 221-246, 1980.
- [8] H. M. Srivastava, and J.L. Brenner, Bounds for Jacobi and related polynomials derivable by matrix methods, *J. Approx. Theory* 12,372-377, 1974.
- [9] H. M. Srivastava, and S. Panda, Some general theorem on generating functions and their applications, *Appl. Math. Lett.* 1, 1988.
- [10] H. M. Srivastava, and H. L. Manocha, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane, and Toronto, 1984.
- [11] H. M. Srivastava, and R. Panda, On the unified presentation of certain classical polynomials, *Boll. Un. Hat Ital.* (4) 12, 306-314, 1975.
- [12] H. M. Srivastava, and B. S. Popov, A class of generating functions for the Jacobi and related polynomials, *Studies in Appl. Math.* 78,177-182, 1988.
- [13] G. Szego, *Orthogonal Polynomials*, Fourth ed. (Amer. Math. Soc. Colloq. Publ. 23), Amer. Math Soc., Providence, Rhode Island, 1975.
- [14] L. N. Mishra, R. K. Yadav, and Deepmala, A note on ordinary hypergeometric series and Bailey's Transform, *Journal of Fractional Calculus and Applications*, Vol. 11, No. 1, 182-187, 2020.
- [15] S. P. Singh, L. N. Mishra, and V. Yadav, Elliptic Well-Poised Bailey Lemma and its Applications, *Journal of Fractional Calculus and Applications*, Vol. 10, No. 2, 31-39, 2019.
- [16] V. N. Mishra, S. N. Singh, S.P. Singh, and V. Yadav, On q-series and continued fractions, *Cogent Mathematics*, 3: 1240414, 1-7, 2016.
- [17] L. N. Mishra, On existence and behavior of solutions to some nonlinear integral equations with applications, Ph.D. Thesis, National Institute of Technology, Silchar 788 010, Assam, India, 2017.
- [18] V. N. Mishra, Some Problems on Approximations of Functions in Banach Spaces, Ph.D. Thesis, Indian Institute of Technology, Roorkee 247 667, Uttarakhand, India, 2007.
- [19] V. N. Mishra, and L. N. Mishra, Trigonometric Approximation of Signals (Functions) in  $L_p$ -norm, *International Journal of Contemporary Mathematical Sciences*, Vol. 7, no. 19, 909-918, 2012.
- [20] R. Dubey, Deepmala, and V. N. Mishra, Higher-order symmetric duality in nondifferentiable multiobjective fractional programming problem over cone constraints, *Stat., Optim. Inf. Comput.*, Vol. 8, 187–205. DOI: 10.19139/soic-2310-5070-601, 2020.
- [21] S. Pandey, Approximation by some Linear Positive Operators using one and two parameters quantum algebra, Ph.D. Thesis, Sardar Vallabhbhai National Institute of Technology, Surat 395 007, Gujarat, India, 2019.

- [22] V. N. Mishra, and S. Pandey, On  $(p, q)$  Baskakov-Durrmeyer-Stancu Operators, Advances in Applied Clifford Algebras, Vol. 27, Issue 2, 1633-1646. DOI: 10.1007/s00006-016-0738-y, 2017.
- [23] V. N. Mishra, and S. Pandey, Certain Modifications of  $(p, q)$ - Szász -Mirakyan Operator, Azerbaijan Journal of Mathematics, Vol. No. 9, issue 2, 2019.
- [24] V. N. Mishra, S. Pandey, and I.A. Khan, On a modification of Dunkl generalization of Szász Operators via  $q$ -calculus, European Journal of Pure and Applied Mathematics, Vol. no. 10, issue no 5, 1067-1077, 2017.
- [25] V.N. Mishra, R.N. Mohapatra, and P. Sharma, On approximation properties of Baskakov-Szász-Stancu operators using Hypergeometric representation, Applied Math. Comp., Vol. 294, 77-86. doi: 10.1016/j.amc.2016.08.049, 2017.
- [26] V.N. Mishra, H.H. Khan, K. Khatri, and L.N. Mishra, Hypergeometric Representation for Baskakov-Durrmeyer-Stancu Type Operators, Bulletin of Mathematical Analysis and Applications, Volume 5 Issue 3, 18-26, 2013.
- [27] V.N. Mishra, and P. Sharma, Direct estimates for Durrmeyer-Baskakov-Stancu type operators using Hypergeometric representation, Journal of Fractional Calculus and Applications, Vol. 6, 2, 1-10, 2015.
- [28] E. Mittal, S. Joshi, R.M. Pandey, and V.N. Mishra, Fractional integral and integral transform formulae using generalized Appell hypergeometric function, Nonlinear Science Letters A, Vol. 8, No. 2, 221-227, 2017.

LAKSHMI NARAYAN MISHRA

DEPARTMENT OF MATHEMATICS, SCHOOL OF ADVANCED SCIENCES, VELLORE INSTITUTE OF TECHNOLOGY (VIT) UNIVERSITY, VELLORE 632 014, TAMIL NADU, INDIA

*Email address:* lakshminarayanmishra04@gmail.com, l.n.mishra@yahoo.co.in

RAKESH KUMAR SINGH

DEPARTMENT OF MATHEMATICS, NETAJI SUBHAS INSTITUTE OF TECHNOLOGY, BIHTA 801 118, PATNA, INDIA

*Email address:* rakeshkumarsingh117@gmail.com

SHIKHA PANDEY

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCES AND LANGUAGES, VIT-AP UNIVERSITY, AMARAVATI 522 237, ANDHRA PRADESH, INDIA

*Email address:* sp1486@gmail.com, shikha.pandey@vitap.ac.in